

Draft Report

Ecological Assessment for Stage Significant Values: Merrimu Precinct Structure Plan, Victoria

Prepared for

Bacchus Marsh Developments Pty Ltd

February 2019



Ecology and Heritage Partners Pty Ltd

DOCUMENT CONTROL

Assessment	Ecological Assessment for Stage Significant Values
Address	Merrimu Precinct Structure Plan, Victoria
Project number	11720
Project manager	Shannon LeBel (Senior Ecologist)
Report reviewer	Shannon LeBel (Senior Ecologist)
Mapping	Monique Elsley (GIS Co-ordinator)
File name	11720_EHP_EA_Vic_MerrimuPSP_Draft_01022019
Client	Bacchus Marsh Developments Pty Ltd
Bioregion	Victorian Volcanic Plain and Central Victorian Uplands
CMA	Port Philip and Westernport
Council	Moorabool Shire Council

Report versions	Comments	Comments updated by	Date submitted
Draft	Submitted to client for review		01/02/2019

Acknowledgements

We thank the following people for their contribution to the project:

- Yvonne Bartonek and Nick Parthemos (Bacchus Marsh Developments) for project and site information;
- Tim Peggie (Ethos Urban) for project information;
- The landowners who provided access to the study area;
- The Victorian Department of Environment, Land, Water and Planning and the Commonwealth Department of the Environment and Energy for access to ecological databases.

Copyright © Ecology and Heritage Partners Pty Ltd

This document is subject to copyright and may only be used for the purposes for which it was commissioned. The use or copying of this document in whole or part without the permission of Ecology and Heritage Partners Pty Ltd is an infringement of copyright.

Disclaimer

Although Ecology and Heritage Partners Pty Ltd have taken all the necessary steps to ensure that an accurate document has been prepared, the company accepts no liability for any damages or loss incurred as a result of reliance placed upon the report and its contents.

SUMMARY

Introduction

Ecology and Heritage Partners Pty Ltd was commissioned by Bacchus Marsh Developments Pty Ltd to conduct a detailed ecological assessment within 16 properties that are landholdings that Bacchus Marsh Developments Pty Ltd have an interest in.

The properties are within an area identified for potential future urban development as part of the expansion of Bacchus Marsh, and Moorabool Council and the Victorian Planning Authority (VPA) have jointly prepared the draft Bacchus Marsh Urban Growth Framework (UGF).

This report summarises the extent of State-significant ecological values present within the Bacchus Marsh Developments landholdings.

Methods

Ecological Assessment

A series of field assessments were undertaken between 15 August and 8 December 2017, and 3 July 2018 to obtain information on terrestrial flora and fauna values within the study area. A habitat hectare assessment was undertaken in conjunction with the flora survey. Vegetation within the study area was assessed according to the habitat hectare methodology, which is described in the Vegetation Quality Assessment Manual.

Targeted Spiny Rice-flower Survey

A targeted survey for the nationally significant Spiny Rice-flower *Pimelea spinescens* subsp. *spinescens* was undertaken on 17, 21, 24 and 31 August, and 4 and 5 September 2017, and 3 July 2018, with the surveys on each of the respective days undertaken by up to four qualified ecologists familiar with the target species.

Areas identified as supporting suitable habitat were traversed, with surveys conducted along transect lines approximately five metres apart, or as dictated by the density of existing grasses and weeds. The location of all plants was recorded during the survey with a handheld GPS (accuracy of +/- 3 metres).

Results

Flora

Remnant native vegetation in the study area is representative of four EVCs: *Low Rainfall* Plains Grassland (EVC 132_63), Grassy Woodland (EVC 175), Rocky Chenopod Woodland (EVC 64) and Plains Grassy Wetland (EVC 125).

A summary of the breakdown of native vegetation recorded in the study area is provided below (Table S1).

TableS1. Summary of native vegetation within the study area

Native Vegetation	Properties 1-15	Property 16	Total
Scattered Trees	60	0	60
Grassy Woodland	4.776	0	4.776
Plains Grassland	19.358	15.095	34.453
Plains Grassy Wetland	0.069	0	0.069
Rocky Chenopod Woodland	0.371	27.799	28.170
Current Wetland	5.323	0	5.323
Remnant Patch (total)	29.897	42.894	72.791

The State significant Spiny Rice-flower, Fragrant Saltbush *Rhagodia parabolica*, Black Roly-poly *Sclerolaena muricata* var. *muricata*, Slender Bindweed *Convolvulus angustissimus* subsp. *omnigracilis*, Melbourne Yellow-gum *Eucalyptus leucoxylon* subsp. *connata* and Bacchus Marsh Wattle *Acacia rostriformis* were recorded within the study area.

Targeted surveys have been undertaken for Spiny Rice-flower at an appropriate time of year within suitable habitat to ascertain its presence within the study area (Section 3.5).

Based on habitat condition and the proximity of previous records, there is suitable habitat within the study area for the State-significant Buloke *Allocasuarina luehmannii*, Small Scurf-pea *Cullen parvum*, Arching Flax-lily *Dianella* sp. aff. *longifolia* (Benambra), and Austral Tobacco *Nicotiana suaveolens*. However, these species were not observed during the ecological assessments.

Fauna

Targeted surveys recorded approximately 50 hectares of confirmed habitat for the Golden Sun Moth *Synemon plana*.

No Striped Legless Lizard *Delma impar* were recorded during the targeted surveys.

Based on habitat condition, and the proximity of previous records, there is also potential habitat within the study area for the State-significant Speckled Warbler *Chthonicola sagittatus*, Barking Owl *Ninox connivens* and Crested Bellbird *Oreoica gutturalis* as well as the Regionally significant Fat-tailed Dunnart and Spotted Harrier *Circus assimilis*.

Communities

One remnant of habitat zone PG4 and all of PG7, PG8 and PG9 meet the description of the FFG Act listed Western (Basalt) Plains Grassland Community ecological community.

A total of 15.835 hectares of the community was recorded within the study area, with 14.456 hectares located in Property 16, and 1.379 hectares situated in Property 11. All areas of this community are proposed to be retained.

Habitat zones RCW3, RCW5 and RCW5 meet the description of the State significant Rocky Chenopod Open Scrub Community ecological community. A total of 27.799 hectares of the Rocky Chenopod Open Scrub Community ecological community is present. All areas of this community are proposed to be retained.

Legislative and Policy Implications

Flora and Fauna Guarantee Act 1988 (FFG Act - Victoria)

There is suitable habitat within the study area for several species listed or protected under the FFG Act (Section 3.4). However, the study area is privately owned, as such a permit under the FFG Act is not required, unless impacts to FFG Act listed matters are impacted on public land (i.e. road reserves). No FFG Act listed communities will be impacted by the proposed development. If required, the proponent should allow up to six weeks to obtain a FFG Act permit through DELWP.

Environment Effects Act 1978

A total of 12.069 hectares of native vegetation is to be impacted as part of the development, which consists of 5.3201 hectares of 'Current Wetland', and a total of 6.7489 hectares of native vegetation as mapped during the field assessments.

A hydrological assessment of the wetland concluded that a more suitable representation of the area likely to support a shallow ephemeral wetland is approximately 2 hectares in size (Water Technology 2019).

Based on a review of ecological impacts associated with the proposed development, it is Ecology and Heritage Partners' position that a referral under the EES Act is not required based on ecological impacts alone.

DRAFT

Table S1. Application requirements for a permit to remove native vegetation (*Victoria Planning Provisions Clause 52.17 -3; DELWP 2017a*)

No.	Application Requirement	Response
Application requirements under the Detailed Assessment Pathway		
1	Information about the native vegetation to be removed, including: <ul style="list-style-type: none"> The assessment pathway and reason for the assessment pathway. A description of the native vegetation to be removed: Maps showing the native vegetation and property in context: The offset requirement that will apply if the native vegetation is approved to be removed. 	Details provided in Section 3 and NVR report in Appendix 4.
2	Topographic and land information relating to the native vegetation to be removed, showing ridges, crests and hilltops, wetlands and waterways, slopes of more than 20 percent, drainage lines, low lying areas, saline discharge areas, and areas of existing erosion, as appropriate.	Details provided in Section 1.3 and Figure 2.
3	Recent, dated photographs of the native vegetation to be removed.	Details provided in Section 3.
4	Details of any other native vegetation approved to be removed, or that was removed without the required approvals, on the same property or on contiguous land in the same ownership as the applicant, in the five year period before the application for a permit is lodged.	Not Applicable.
5	An avoid and minimise statement. The statement describes any efforts to avoid the removal of and minimise the impacts on the biodiversity and other values of native vegetation, and how these efforts focussed on areas of native vegetation that have the most value.	Details provided in Section 5.1.
6	A copy of any Property Vegetation Plan contained within an agreement made pursuant to section 69 of the <i>Conservation, Forests and Lands Act 1987</i> that applies to the native vegetation to be removed.	Not applicable.
7	Where the removal of native vegetation is to create defensible space, a written statement explaining why the removal of native vegetation is necessary. This statement must have regard to other available bushfire risk mitigation measures. This statement is not required when the creation of defensible space is in conjunction with an application under the Bushfire Management Overlay.	Not applicable.
8	If the application is under Clause 52.16, a statement that explains how the proposal responds to the Native Vegetation Precinct Plan considerations at decision guideline 8.	Not applicable.
9	An offset statement providing evidence that an offset that meets the offset requirements for the native vegetation to be removed has been identified and can be secured in accordance with the Guidelines.	Details provided in Section 5.3
10	A site assessment report of the native vegetation to be removed, including: <ul style="list-style-type: none"> A habitat hectare assessment of any patches of native vegetation, including the condition, extent (in hectares), Ecological Vegetation Class and bioregional conservation status. The location, number, circumference (in centimetres measured at 1.3 metres above ground level) and species of any large trees within patches. The location, number, circumference (in centimetres measured at 1.3 metres above ground level) and species of any scattered trees, and whether each tree is small or large. 	See Section 3.3; Appendix 2.3.
11	Information about impacts on rare or threatened species habitat, including: <ul style="list-style-type: none"> The relevant section of the Habitat importance map for each rare or 	Details provided in Section 3.4 and Appendix 4.

No.	Application Requirement	Response
	<p>threatened species requiring a species offset.</p> <ul style="list-style-type: none"> • For each rare or threatened species that the native vegetation to be removed is habitat for, according to the Habitat importance maps: <ul style="list-style-type: none"> - the species' conservation status - the proportional impact of the removal of native vegetation on the total habitat for that species - whether their habitats are highly localised habitats, dispersed habitats, or important areas of habitat within a dispersed species habitat 	

DRAFT

CONTENTS

1	INTRODUCTION	9
1.1	Background	9
1.2	Objectives	9
1.3	Study Area.....	9
2	METHODS	11
2.1	Desktop Assessment	11
2.2	Field Assessment.....	11
2.3	Removal, Destruction or Lopping of Native Vegetation (the Guidelines)	13
2.4	Assessment Qualifications and Limitations	13
3	RESULTS	14
3.1	Vegetation Condition	14
3.2	Fauna Habitat.....	21
3.3	Removal of Native Vegetation (the Guidelines).....	21
3.4	Significance Assessment	23
3.5	Targeted Spiny Rice-flower Survey	27
4	LEGISLATIVE AND POLICY IMPLICATIONS	28
4.1	<i>Flora and Fauna Guarantee Act 1988</i> (Victoria).....	28
4.2	<i>Environment Effects Act 1978</i>	28
5	MITIGATION MEASURES	33
5.1	Avoidance and Minimisation Statement.....	33
5.2	Best Practice Mitigation Measures	33
5.3	Offset Impacts.....	34
6	FURTHER REQUIREMENTS.....	36
	REFERENCES.....	37
	FIGURES	40
	APPENDICES	48
	APPENDIX 1.....	49
	APPENDIX 2 - FLORA.....	55
	APPENDIX 3 - FAUNA.....	68
	APPENDIX 4 - NATIVE VEGETATION REPORTS	73

1 INTRODUCTION

1.1 Background

Ecology and Heritage Partners Pty Ltd was commissioned by Bacchus Marsh Developments Pty Ltd to conduct an Ecological Assessment for Stage Significant Values within the 17 properties that are proposed to comprise the Merrimu Precinct Structure Plan, Victoria.

Ecology and Heritage Partners understand that Bacchus Marsh Developments Pty Ltd has acquired interests in the properties, which are currently used for agriculture. The properties are within an area identified for potential future urban development as part of the expansion of Bacchus Marsh, and Moorabool Council and the Victorian Planning Authority (VPA) have jointly prepared the draft Bacchus Marsh Urban Growth Framework (UGF).

This report is a subset of the broader Ecological Assessment prepared for Bacchus Marsh Developments (Ecology and Heritage Partners 2018a), and only describes values of State significance recorded within the Bacchus Marsh Developments landholdings (Figure 1). As such, the majority of the methodology and legislative implications are not repeated herein, with only the results of the assessments as they relate to State significant values provided in this report.

1.2 Objectives

The purpose of the assessment was to identify the extent and type of remnant native vegetation present within the properties of which Bacchus Marsh Developments has an interest in, and to determine the presence, or otherwise, of State significant flora and fauna species and/or ecological communities. This report presents the results of the ecological assessments and discusses the potential ecological and legislative implications associated with any future proposed development.

Where areas of remnant vegetation were present, the following tasks were completed to address requirements under the 'Guidelines for the removal, destruction or lopping of native vegetation (The Guidelines) (DELWP 2017a):

- A habitat hectare assessment of any areas of remnant native vegetation within the study area;
- Quantify the presence of scattered trees and Large Trees in patches (LOTs) and habitat for rare or threatened species that may be impacted as a result of the proposed development.

1.3 Study Area

The study area covers approximately 435 hectares and is comprised of 17 properties bound by Gisborne Road to the west, and Bences Road to the east (Table 1) approximately 50 kilometres north-west of Melbourne's CBD (Figure 1). It should be noted that Property 16 is ultimately proposed to be secured and managed as an offset site and will not be subject to any proposed development.

Table 1. Properties within the study area

Property Number *	Address	Title details
1	2621 Gisborne Road	Proposed lots 1 and 2 on PS724534Y
2	Gisborne Road	Lot 1 PS124024
3	146 Bences Road	PS124024
4a	2677 Gisborne Road	Lot 1 TP578035R
4b	152 Bences Road	Lot 1 TP159956
5	Gisborne Road	Lots 1,2,3&4 TP567257J
6	Buckleys Road	Lot 1 on TP958042C
7	268 Bences Road	Lot 1 PS125141
8	139 O'Connell Road	Lots 1&2 TP408175C
9	332 Bences Road	Lot 2 PS125141
10	372 Bences Road	Lot 2 PS432900C
11	376 Bences Road	Lot 2 PS411883S
12	Lerderderg Park Road	Lot 1 TP97760S
13	Lerderderg Park Road	Lot 1 TP111405 (part)
14	345 Bences Road	Lot 2 PS139808
15	295 Bences Road	Lot 1 LP139808
16*	289 Bences Road	Allot E, Sec 18\PP3095

Note. See Figure 2 for location of properties within the study area. * Property 16 is not proposed for development and is proposed to be secured and protected as an offset site.

The land within and surrounding the study area predominantly supports agricultural activities in the form of grazing, cropping, market gardens, orchards, and vineyards. Two operating quarries are located immediately opposite the study area on the west of Gisborne Road, while the Long Forest Flora and Fauna Reserve is located to the east of Bences road in close proximity to the study area (Figure 1).

The study area is generally flat, with several escarpments located to the west and south of the study area. The headwaters of several designated waterways commence within the study area and follow the escarpments into lower lying areas to the east and west.

Erosion is evident throughout all observed escarpments and has resulted in a shallow soil profile at both the top and mid-slope of these the escarpments. The location of waterways, escarpments, steep slopes and erosion within the study area is shown in Figure 1 and Figure 2.

According to the Department of Environment, Land, Water and Planning (DELWP) Native Vegetation Information Management (NVIM) Tool (DELWP 2019a), the study area occurs within the Victorian Volcanic Plain and Central Victorian Uplands bioregions. It is located within the jurisdiction of the Port Philip and Westernport Catchment Management Authority (CMA) and the Moorabool Shire Council municipality. Section 0 discusses zoning and overlays relevant to the study area.

2 METHODS

2.1 Desktop Assessment

A detailed methodology used to inform the desktop assessment is included in Ecology and Heritage Partners (2018a).

2.2 Field Assessment

2.2.1 Flora Assessment

The flora assessment was undertaken on 15 -17 August, 10 September, 25 October, 8 December 2017 and 3 July 2018 to obtain information on flora values within the study area. The study area was walked and/or driven, with all observed vascular flora species recorded, any significant records mapped, and the overall condition of vegetation noted. Remnant vegetation in the local area was also investigated to assist in determining the pre-European vegetation within the study area. EVCs were determined with reference to DELWP pre-1750 and extant EVC mapping and their published descriptions (DELWP 2019c). The significance assessment criteria of taxa and vegetation communities are presented in Appendix 1.

Where remnant vegetation was identified a habitat hectare assessment was undertaken following methodology described in the Vegetation Quality Assessment Manual (DSE 2004).

2.2.1.1 Spiny Rice-flower Targeted Survey

A targeted survey for the State significant Spiny Rice-flower *Pimelea spinescens* subsp. *spinescens* was undertaken on 17, 21, 24 and 31 August, and 4 and 5 September 2017, and 3 July 2018 with the survey on each of the respective days undertaken by up to four qualified ecologists familiar with the target species. Areas identified as supporting suitable habitat (Properties 4b, 5, 9, 10, 11, 15 and 16) were traversed, with surveys conducted along transect lines approximately five metres apart, or as dictated by the density of existing grasses and weeds. The location of all plants was recorded during the survey with a handheld GPS (accuracy of +/- 3 metres).

The survey methodology adhered to the survey guidelines for Spiny Rice-flower outlined in the Biodiversity Precinct Structure Planning Kit (DSE 2010a) and in the Significant Impact Guidelines for the species (DEWHA 2009). A summary of the survey effort compared with the survey guidelines is provided in Table 2.

Spiny Rice-flower is a perennial sub-shrub listed as threatened under the Victorian FFG Act, and as endangered under the Advisory List of Rare and Threatened Plants in Victoria (DEPI 2014). The species is endemic to Victoria and is found between the south-west and north-central parts of the State. It occurs in grassy EVC such as Plains Grassland (EVC 132), Plains Grassy Woodland (EVC 55), Plains Woodland (EVC 803) and Plains Grassland/Grassy Woodland Mosaic (EVC 897) (DEWHA 2009). Spiny Rice-flower is typically found in small populations (<500 individuals).

The species is slow-growing and reaches up to 30 cm in height (Plate 1; Plate 2). Plants are mostly dioecious (male and female flowers on separate plants) but some plants are monoecious (male and female flower on same plant). It bears small yellow flowers between April and August (DEWHA 2009).

Table 2. Survey effort compared with the Biodiversity Precinct Structure Planning Kit (DSE 2010a) and the Significant Impact Guidelines for the species (DEWHA 2009).

Survey Guidelines	Comment
Targeted surveys should be done by people familiar with recognising the subspecies.	Yes. Surveys were completed by assessors familiar with the appearance and ecology of the subspecies.
Multiple surveys may be required to identify the species and provide adequate survey effort.	Given that the species was known to be flowering at the time of the assessments, and biomass was generally low across areas of suitable habitat, specimens were easily identifiable, a single survey effort across most of the properties was considered appropriate to accurately record the species. Multiple surveys were undertaken in Property 11 and 16 where large populations were identified.
Surveys should not be conducted for at least six months after fires and for at least three months after the cessation of grazing (DEWHA Survey Guidelines).	Yes. The assessors are not aware of any fires or grazing within the specified timeframes.
Survey Spiny Rice-flower between April and August (easily overlooked when not in flower).	Yes. The assessments were conducted within the flowering period for the species by ecologists familiar with the species in and out of flower. Given the survey effort within areas of suitable habitat, there is reasonable assurance that individuals were not overlooked.
The targeted survey effort should be directed to all potential habitat areas i.e. remnant grassland including degraded grassland.	Yes. The entire study area was visually surveyed and traversed in linear transects (i.e. targeted survey areas).
Walk through transects at less than 5m grid intervals are required for all potential habitat.	Yes. Transects of five metres apart were utilised throughout the entire targeted survey areas.
Record the number of plants per land parcel.	Yes. Any observed plants were recorded.



Plate 1. Spiny Rice-flower within the study area (Ecology and Heritage Partners Pty Ltd 2017).



Plate 2. Spiny Rice-flower within the study area (Ecology and Heritage Partners Pty Ltd 2017).

2.2.2 Fauna Assessment

A fauna assessment was undertaken on 15 and 16 August 2017 to obtain information on terrestrial fauna values within the study area. The study area was visually assessed and active searching under and around ground debris for reptiles, frogs and small mammals was undertaken. Binoculars were also used to scan the area for birds, and observers listened for calls and searched for other signs of fauna such as nests, remains of dead animals, droppings and footprints. Potential habitat for fauna was assessed, with a particular emphasis on habitats that may provide shelter, food or other resources for significant species.

2.3 Removal, Destruction or Lopping of Native Vegetation (the Guidelines)

Assessment methodology and background relating to the 'Guidelines for the removal, destruction or lopping of native vegetation' (the Guidelines) (DELWP 2017a) is provided in Ecology and Heritage Partners (2018a).

2.4 Assessment Qualifications and Limitations

Data and information held within the ecological databases and mapping programs reviewed in the desktop assessment (e.g. VBA, PMST, Online databases and maps etc.) are unlikely to represent all flora and fauna observations within, and surrounding, the study area. It is therefore important to acknowledge that a lack of documented records does not necessarily indicate that a species or community is absent.

Ecological values identified on site are recorded using a hand-held GPS or tablet with an accuracy of +/-3 metres. This level of accuracy is considered adequate to provide an accurate assessment of the ecological values present within the study area; however, this data should not be used for detailed surveying purposes.

The 'snap shot' nature of a standard biodiversity assessment means that migratory, transitory or uncommon fauna species may have been absent from typically occupied habitats at the time of the field assessment. In addition, annual or cryptic flora species such as those that persist via underground tubers may also be absent. Targeted flora or fauna surveys were not undertaken, as this was beyond the preliminary scope of the project. Nevertheless, the terrestrial flora and fauna data collected during the field assessment and information obtained from relevant desktop sources is considered adequate to provide an accurate assessment of the ecological values present within the study area.

Where appropriate, a precautionary approach has been adopted in the discussion of implications. That is, where insufficient evidence is available on the occurrence or likelihood of occurrence of a species, it is assumed that it could be in an area of habitat, if suitable, and the implications under legislation and policy are considered accordingly.

It is noted that within this report, the identification of the Waxy Yellow-gum *Eucalyptus leucoxylon* subsp. *pruinosa* included in Ecology and Heritage Partners (2018a) has been re-assessed and reclassified as Melbourne Yellow-gum *Eucalyptus leucoxylon* subsp. *connata* based on recent field assessments conducted within Property 16 and other nearby adjacent properties undertaken in December 2018 (Ecology and Heritage Partners *in prep*).

3 RESULTS

3.1 Vegetation Condition

3.1.1 Remnant Patches

Remnant native vegetation in the study area is representative of four EVCs: *Low Rainfall* Plains Grassland (EVC 132_63), Grassy Woodland (EVC 175), Rocky Chenopod Woodland (EVC 64) and Plains Grassy Wetland (EVC 125). The presence of these EVCs is generally consistent with the modelled pre-1750s native vegetation mapping (DELWP 2019a).

The remainder of the study area comprises introduced and planted vegetation, present as crop, pasture, windrows and ornamental plantings. Specific details relating to observed EVCs are provided below.

3.1.1.1 Plains Grassland

Low-rainfall Plains Grassland (EVC 132_63) typically consists of treeless vegetation mostly less than one metre in height, and dominated by a mixture of grasses and herbs. This EVC usually occupies cracking basalt soils prone to seasonal waterlogging in areas receiving less than 500 millimetres of annual rainfall (DELWP 2019c).

Plains Grassland was recorded along the north-west boundary, and in scattered patches to the south and east of the study area (Figure 2). Dominant native grasses recorded throughout most patches included Spurred Spear-grass *Austrostipa gibbosa*, Rough Spear-grass *Austrostipa scabra* subsp. *falcata*, Common Wallaby-grass *Rytidosperma caespitosa*, Bristly Wallaby-grass *Rytidosperma setaceum*, and Knead Wallaby-grass *Rytidosperma geniculatum* (Plate 3). Commonly observed shrubs and herbs within this vegetation type comprised Berry Saltbush *Atriplex semibaccata*, Sheep's Burr *Acaena echinata*, Wingless Bluebush *Maireana enchylaenoides*, Nodding Saltbush *Einadia nutans*, Ruby Saltbush *Enchylaena tomentosa* var. *tomentosa*, Native Flax *Linum marginale* and occasional specimens of Lemon Beauty-heads *Calocephalus citreus*, Fuzzy New Holland Daisy *Vittadinia cuneata*, and Golden Billy-buttons *Pycnosorus chrysanthes* (Plate 4).

The State significant Slender Bindweed *Convolvulus angustissimus* subsp. *omnigracilis*, Black Roly-poly *Sclerolaena muricata* var. *muricata*, Spiny Rice-flower *Pimelea spinescens* subsp. *spinescens* and Fragrant Saltbush *Rhagodia parabolica* were recorded within several Plains Grassland remnants (Figure 2; Figure 3).

A total of 10 habitat zones comprising 34.453 hectares were recorded within the study area (PG1 – PG10) (Figure 2), with habitat zones differing in quality predominantly due to the diversity and/or of native species present, and the type and extent of weeds present in the habitat zone (Appendix 2.3). A total of 15.095 hectares of Plains Grassland is located in Property 16.

Some remnants of habitat zone PG4 (Figure 2b), and all of PG8, PG9 and PG10 (Figure 2c) meet the description of the State significant *Western (Basalt) Plains Grassland* ecological community.

Patches PG8 and PG9 were of the highest quality, were contiguous with each other and other larger remnants of vegetation in Property 16 and supported high native species diversity.



Plate 3. Plains Grassland (PG2) within the study area (Ecology and Heritage Partners Pty Ltd 17/08/2017).



Plate 4. Fuzzy New Holland Daisy-dominated Plains Grassland (PG8) within the study area (Ecology and Heritage Partners Pty Ltd 10/09/2017).

Lower quality remnants were located elsewhere throughout the study area, and due to former or ongoing land practices, exhibited a lower species diversity, with PG2, PG3 and PG6 often being defined by only one or two native species, and high cover of exotic flora.

Exotic flora was dominant throughout most areas within and adjacent to Plains Grassland vegetation. The most commonly observed weeds were the declared Victorian noxious weeds African Box-thorn *Lycium ferocissimum*, Artichoke Thistle *Cynara cardunculus*, Horehound *Marrubium vulgare* and Serrated Tussock *Nassella trichotoma*. Other common environmental weeds present throughout included Galenia *pubescens*, Cape Weed *Arctotheca calendula*, Wild Turnip *Brassica* spp., Perennial Rye-grass *Lolium perenne*, Barley *Hordeum* spp., Rat's-tail Fescue *Vulpia myuros*, Ribwort *Plantago lanceolata* and Soft Brome *Bromus hordeaceus* (Plate 5; Plate 6).



Plate 5. Galenia and African Box-thorn within the study area (Ecology and Heritage Partners Pty Ltd 15/08/2017).



Plate 6. Serrated Tussock-dominated grassland (Ecology and Heritage Partners Pty Ltd 25/10/2017).

3.1.1.2 Grassy Woodland

Grassy Woodland is described as a variable open eucalypt woodland over a diverse ground layer of grasses and herbs, with a sparse shrub component. The EVC usually occurs on sites with moderate fertility over a range of geologies, often on undulating hillsides or slopes (DELWP 2019c).

Within the study area, Grassy Woodland was recorded in several small, scattered remnants adjacent to Gisborne Road, along with one large remnant immediately north of O'Connell Road (Figure 2).

The overstorey was predominantly comprised of Grey Box *Eucalyptus microcarpa*, with occasional specimens of Yellow Box *Eucalyptus melliodora*, and the State significant Melbourne Yellow Gum *Eucalyptus leucoxylon* subsp. *connata* (Figure 2b and 2c).

The understory was in poor condition in all habitat zones, with only occasional occurrences of native grasses and shrubs present. The State significant Fragrant Saltbush was relatively common within and adjacent to several patches of Grassy Woodland. However, the dominant understory species comprised Galenia and the noxious weeds African Box-thorn and Serrated Tussock (Plate 7; Plate 8).

A total of four habitat zones were recorded within the study area (GW1 – GW4) (Figure 2), comprising an area of 5.486 hectares, with habitat zones mostly defined by the number of Large Old Trees present, and the cover of weeds in the understory (Appendix 2.3).



Plate 7. Grassy Woodland (GW₁) within the study area (Ecology and Heritage Partners Pty Ltd 25/10/2017).



Plate 8. Grassy Woodland (GW₄) within the study area (Ecology and Heritage Partners Pty Ltd 16/08/2017).

3.1.1.3 Rocky Chenopod Woodland

Rocky Chenopod Woodland is a low open eucalypt woodland (often in mallee-form) with an understory dominated by chenopod (saltbush) species, with scattered grasses and herbs (DELWP 2019c).

Rocky Chenopod Woodland was recorded in small patches near north-west boundary adjacent to Gisborne Road, as well as in a large remnant in the north-eastern half of the study area in Property 16 (Figure 2c). The overstorey of this EVC was co-dominated by Grey Box and Bull Mallee *Eucalyptus behriana*, with the occasional Melbourne Yellow Gum specimen also present. The understory was generally sparse, and comprised Fragrant Saltbush, Ruby Saltbush, Moonah *Melaleuca lanceolata*, Gold-dust Wattle *Acacia acinacea*, Variable Groundsel *Senecio pinnatifolius* and Saloop *Einadia hastata* (Plate 9 and 10).

Weed cover was high in habitat zones RCW1, RCW2 and RCW5, with African Box-thorn and Galenia dominating the understory in these habitat zones (Plate 11; Plate 12). Zones RCW3 and RCW4 were relatively weed free (Plate 9; Plate 10).

Five habitat zones (RCW1 – RCW5) were recorded comprising a total of 28.121 hectares, of which 27.788 hectares is situated in Property 16. Habitat zones were predominantly differentiated due to the cover of weeds and number of woody species exhibiting recruitment (Appendix 2.3).

Habitat zones RCW3, RCW4 and RCW5 in Property 16 meet the description of the State significant *Rocky Chenopod Open Scrub* ecological community.



Plate 9. Rocky Chenopod Woodland (RCW₃) within the study area (Ecology and Heritage Partners Pty Ltd 08/12/2017).



Plate 10. Rocky Chenopod Woodland (RCW₃) within the study area (Ecology and Heritage Partners Pty Ltd 08/12/2017).



Plate 11. High cover of African Box-thorn within RCW₅ (Ecology and Heritage Partners Pty Ltd 08/12/2017).



Plate 12. High cover of African Box-thorn within RCW₅ (Ecology and Heritage Partners Pty Ltd 08/12/2017).

3.1.1.4 Plains Grassy Wetland

Plains Grassy Wetland is usually treeless, although a sparse shrub component may be present. The ground cover is usually dominated by grasses and small sedges and herbs. The vegetation is typically species-rich on the outer verges but is usually species-poor in the wetter central areas (DELWP 2019c).

One patch of Plains Grassy Wetland (PGWe1) was recorded around an artificial water body along a designated waterway within Property 5, and was comprised of Cumbungi *Typha* spp., Common Spike-sedge *Eleocharis acuta*, Joint-leaf Rush *Juncus holoschoenus* and Pale Rush *Juncus pallidus* (Plate 13; Plate 14).

The exotic species Drain Flat-sedge *Cyperus eragrostis*, Lesser Quaking-grass *Briza minor* and Yorkshire Fog *Holcus lanatus* were common in this habitat zone.



Plate 13. PGWe1 within the study area (Ecology and Heritage Partners Pty Ltd 15/08/2017).



Plate 14. PGWe1 within the study area (Ecology and Heritage Partners Pty Ltd 15/08/2017).

3.1.2 Scattered Trees

Sixty (60) scattered trees, the majority being Grey Box, with occasional specimens of River Red-gum *Eucalyptus camaldulensis*, Yellow Box, and Messmate *Eucalyptus obliqua* occur throughout the study area with the majority estimated to be at least 200 years old. These trees would once likely have been part of the Grassy Woodland EVC, however the understorey vegetation consists of predominantly introduced species (mainly exotic pasture grasses) and the trees no longer form a patch of native vegetation (Plate 15; Plate 16). Of the 60 scattered trees, a total of 32 are Large Trees (LTs), with 28 Small Trees (STs) (Appendix 2.4).



Plate 15. Two scattered Grey Box within the study area (Ecology and Heritage Partners Pty Ltd 10/09/2017).



Plate 16. Scattered River Red-gum within the study area (Ecology and Heritage Partners Pty Ltd 16/08/2017).

3.1.3 Introduced and Planted Vegetation

3.1.3.1 *Introduced Vegetation*

Areas not supporting remnant native vegetation have a high cover (>80%) of exotic grass species, many of which have been direct-seeded for use as pasture. Scattered native grasses are generally present in these areas, however they did not have the required 25% cover to be considered a remnant patch. Removal of embedded rock has also been undertaken as part of historical agricultural activities throughout much of the study area.

Large areas of the study area have no native vegetation present, and are dominated by cereal crops (Plate 17).

Disturbed areas (not mapped as native vegetation) were mostly dominated by the environmental weeds Galenia, Rat's Tail Fescue, Ribwort, Wild Oat *Avena fatua*, Prairie Grass *Bromus catharticus*, Curled Dock *Rumex crispus*, Black Night-shade *Solanum nigrum*, Sticky Ground-cherry *Physalis hederifolia*, and Onion-grass *Romulea rosea*.

Noxious weeds are present throughout the study area, with common occurrences of Artichoke Thistle, Horehound and Spear Thistle *Cirsium vulgare* along with the Weeds of National Significance (WONS), African Boxthorn, Serrated Tussock, Prickly Pear *Opuntia* spp., Bridal Creeper *Asparagus asparagoides* and Blackberry *Rubus fruticosus* sp. agg. (Plate 18; Plate 19)



Plate 17. Cropped area within the study area (Ecology and Heritage Partners Pty Ltd 16/08/2017).



Plate 18. Artichoke Thistle within the study area (Ecology and Heritage Partners Pty Ltd 16/08/2017).

3.1.3.2 *Planted Vegetation*

Planted vegetation in the study area consists of native and non-native Victorian tree and shrub species. A Sugar Gum *Eucalyptus cladocalyx* plantation is located to the north of the study area (Plate 20).

Aside from Sugar Gum, commonly planted species include Peppercorn *Shoenus molle*, Southern Mahogany *Eucalyptus botryoides*, and specimens of Grey Box and Yellow Gum. Most planted vegetation is situated in windrows, or around dwellings, sheds and laneways. A variety of ornamental shrubs have also been planted around sheds and dwellings.



Plate 19. Artichoke Thistle within the study area (Ecology and Heritage Partners Pty Ltd 08/12/2017).



Plate 20. Planted vegetation within the study area (Ecology and Heritage Partners Pty Ltd 11/02/2017).

3.2 Fauna Habitat

A detailed description of fauna habitat and commonly observed species is included in Ecology and Heritage Partners (2018a).

3.3 Removal of Native Vegetation (the Guidelines)

The study area (Properties 1-16) contains the following extent of native vegetation:

- 60 Scattered Trees, comprising;
 - 32 Large Old Trees; and,
 - 28 Small Trees.
- 72.791 hectares of native vegetation, comprising:
 - 4.776 hectares of the Grassy Woodland EVC;
 - 34.453 hectares of the Plains Grassland EVC;
 - 0.069 hectares of the Plains Grassy Wetland EVC;
 - 28.170 hectares of the Rocky Chenopod Woodland EVC; and,
 - 5.323 hectares of a modelled Current Wetland.

A summary of the breakdown of native vegetation is provided below (Table 3).

Table 3. Summary of native vegetation within the study area

Native Vegetation	Properties 1-15	Property 16	Total
Scattered Trees	60	0	60
Large Trees in patches	40	320	360

Native Vegetation	Properties 1-15	Property 16	Total
Grassy Woodland	4.776	0	4.776
Plains Grassland	19.358	15.095	34.453
Plains Grassy Wetland	0.069	0	0.069
Rocky Chenopod Woodland	0.371	27.799	28.170
Current Wetland	5.323	0	5.323
Remnant Patch (total)	29.897	42.894	72.791

It should be noted that a recent hydrological assessment of the Current Wetland concluded that a more suitable representation of the area likely to support a shallow ephemeral wetland is approximately two hectares in size (Water Technology 2019). An application to reassess the extent of the Current Wetland modelled within Property 5 is currently being prepared for submission to DELWP.

3.3.1 Vegetation proposed to be removed

The extent of native vegetation removal has been determined based on a preliminary impact footprint discussed between Ecology and Heritage Partners and Bacchus Marsh Developments on 25/05/2018.

The study area is within Location category 3, with 12.069 hectares of native vegetation proposed to be removed. As such, the permit application falls under the Detailed assessment pathway. Vegetation impacts are summarised in Table 4.

Condition scores for vegetation proposed to be removed are based on the habitat hectare assessment completed by a qualified vegetation assessor and are provided in Appendix 2.3.

Table 4. Removal of native vegetation (the Guidelines)

Assessment pathway	Detailed
Total Extent	12.069
Large Trees (no.)	0
Location Category	3

3.3.2 Offset Targets

The offset requirement for native vegetation removal is 0.001 General Habitat Units (HUs) and specific offsets for the following species:

- 9.744 species units of habitat for Small Golden Moths *Diuris basaltica*;
- 10.049 species units of habitat for Heath Spear-grass *Austrostipa exilis*;
- 10.108 species units of habitat for Melbourne Yellow-gum *Eucalyptus leucoxylon* subsp. *connata*;
- 9.675 species units of habitat for Basalt Podolepis *Podolepis linearifolia*;
- 10.079 species units of habitat for Bacchus Marsh Wattle *Acacia rostriformis*; and,
- 10.049 species units of habitat for Fragrant Saltbush *Rhagodia parabolica*.

A summary of proposed vegetation losses and associated offset requirements is presented in Table 5 and the Native Vegetation Removal (NVR) Report is presented in Appendix 4.

Table 5. Offset targets

General Offsets Required	0.001 General HUs
Specific Offsets Required	<ul style="list-style-type: none"> • 9.744 species units of habitat for Small Golden Moths <i>Diuris basaltica</i> • 10.049 species units of habitat for Heath Spear-grass <i>Austrostipa exilis</i> • 10.108 species units of habitat for Melbourne Yellow-gum <i>Eucalyptus leucoxylon</i> subsp. <i>connata</i> • 9.675 species units of habitat for Basalt Podolepis <i>Podolepis linearifolia</i> • 10.079 species units of habitat for Bacchus Marsh Wattle <i>Acacia rostriformis</i> • 10.049 species units of habitat for Fragrant Saltbush <i>Rhagodia parabolica</i>
Vicinity (catchment / LGA)	Port Philip and Westernport CMA / Moorabool Shire Council
Minimum SBV	0.800

Note: HU = Habitat Units; SBV – Strategic Biodiversity Value.

3.4 Significance Assessment

3.4.1 Flora

A total of 119 flora species (62 indigenous and 57 non-indigenous or introduced) were recorded within the study area during the field assessment.

The State significant Spiny Rice-flower, Fragrant Saltbush, Black Roly-poly, Slender Bindweed, Melbourne Yellow-gum and Bacchus Marsh Wattle *Acacia rostriformis* were recorded within the study area.

A consolidated list of flora species recorded is provided in Appendix 2.1.

The VBA contains records of 53 State significant flora species previously recorded within 10 kilometres of the study area (DELWP 2019d) (Appendix 2.2; Figure 4). Most records are located within and adjacent to the Long Forest Flora and Fauna Reserve to the east of the study area, the Lerderderg State Park to the north-west of the study area, and existing road reserves within the local area where survey effort has likely been greater (Figure 4).

Based on the condition of remnant vegetation, landscape context and the proximity of previous records, in addition to those observed, the following significant flora species have the potential to occur within the study area (Table 6) (Appendix 2.2).

Table 6. Significant flora species with the potential to occur within the study area.

Common Name	Scientific Name	Habitat
State Significance *		
Small Golden Moths	<i>Diuris basaltica</i>	There are five records of Small Golden Moths recorded in the VBA within the local area, with all located south of Werribee River and Bacchus Marsh township (VBA 2017d). An additional record is located further east near Melton, with another record north-west in Toolern Vale (DELWP 2019d). Small Golden Moth orchids typically grow in herb-rich native grasslands, dominated by

Common Name	Scientific Name	Habitat
		<p>Kangaroo Grass <i>Themeda triandra</i> on heavy basaltic soils, often embedded with basalt boulders, with the known distribution of the species highly restricted (DSE 2010b).</p> <p>Given the absence of Kangaroo Grass-dominated grassland within the study area, general poor condition of habitat (outside of Property 16), high levels of weed invasion, absence of other orchids within the locality, and history of agricultural activities, there is considered to be a low likelihood of occurrence in Properties 1-15, and surveys are not required.</p>
Basalt Peppercross	<i>Lepidium hyssopifolium</i>	<p>Although there are no records within the VBA within 10 kilometres, there is an informal record recorded in the Atlas of Living Australia (ALA) within Property 6 south of Buckleys Road (ALA 2017). This property has been cropped, and the specimen would no longer be present.</p> <p>It is understood that almost all remaining populations of Basalt Peppercross occur in heavily modified, non-natural environments, usually amongst exotic pasture grasses and weed species, sometimes with an overstorey of introduced tree species (DSE 2010c). However, the species appears to rely heavily on favourable microsite conditions, with Basalt Peppercross appearing to only establish in relatively open bare ground where there is limited competition from other plants (both native and introduced species), rather than in areas with thick ground cover (DSE 2010c). As the majority of grassland vegetation (native and non-native_ within Properties 1-15 supports high levels of biomass, with few patches of bare ground present, as well as the lack of other records in close proximity to the study area, there is considered a low likelihood of occurrence within the study area, and targeted surveys are not required.</p> <p>Further, the biodiversity assessment and targeted surveys (for other species) did not note any specimens that meet the description of the species.</p>
Large-head Fireweed	<i>Senecio macrocarpus</i>	<p>There are no known records of Large-headed Fireweed within 10 kilometres of the study area, with the closest known record located approximately 17 kilometres to the east (VBA 2017e). Previous surveys for the species in nearby properties did not record the species (Ecology and Heritage Partners 2013b), and there is considered to be a low likelihood of occurrence that the species occurs in the locality. As such, targeted surveys are not required.</p>
Werribee Blue-box	<i>Eucalyptus baueriana</i> subsp. <i>thalassina</i>	<p>A total of 313 records of Werribee Blue-box have been recorded within 10 kilometres of the study area, with all records located within the Long Forest Flora and Fauna reserve (VBA 2017e; Figure 4).</p> <p>Although there is suitable habitat in Woodland areas throughout the study area, no specimens were observed during the ecological assessment, nor recorded during previous surveys within the study area (BL&A 2010).</p>

Note. * Those species with the highest likelihood of occurrence.

Recommendation

Based on habitat condition, and the proximity of previous records, there is also potential habitat within the study area for the State-significant Buloke *Allocasuarina luehmannii*, Small Scurf-pea *Cullen parvum*, Arching Flax-lily *Dianella* sp. aff. *longifolia* (Benambra), and Austral Tobacco *Nicotiana suaveolens*.

It is noted that the ecological surveys undertaken to date have not recorded these species within the study area, and due to the degraded condition and high weed cover of the majority of properties, the presence of additional State-significant flora is considered to be low.

Targeted surveys for State significant flora would assist to determine their presence within the study area. However, based on existing legislative approvals under the P&E Act, further survey for these species is not currently required to determine offset implications.

3.4.2 Fauna

The VBA contains records of 41 State significant and 13 regionally significant fauna species previously recorded within 10 kilometres of the study area (DELWP 2019d) (Appendix 3.1; Figure 5).

Targeted surveys undertaken by Ecology and Heritage Partners (2018b) confirmed the presence of Golden Sun Moth *Synemon plana* within several properties (Figure 2a-2c). Targeted surveys undertaken for Striped Legless Lizard did not record the species within the study area. A detailed summary of the targeted survey methodology and results is provided in Ecology and Heritage Partners (2018b).

Based on the type and condition of habitats within the study area, landscape context and the proximity of previous records, in addition to those species recorded or surveyed for, the following State significant fauna species have the potential to occur within the study area (Table 7; Appendix 3.1).

Table 7. Significant fauna species with the potential to occur within the study area.

Common Name	Scientific Name	Habitat
State Significance *		
Swift Parrot	<i>Lathamus discolor</i>	Swift Parrot may forage on eucalypts within the study area on occasion. However, the species breeds only in Tasmania and migrates to mainland Australia in autumn and is usually recorded between Stawell in the central west and Wodonga in the north-east. As such the study area is unlikely to provide important or limiting habitat for this species.
Brown Treecreeper	<i>Climacteris picumnus victoriae</i>	There are 75 records of Brown Treecreeper from the local area, with the most recent taken in 2010. Habitat is mainly found in woodland areas within the study area, particularly closer to Long Forest Flora and Fauna Reserve.
Hooded Robin	<i>Melanodryas cucullata cucullata</i>	There are 12 records of Hooded Robin from the local area. Habitat is mainly found in woodland areas within the study area, particularly closer to Long Forest Flora and Fauna Reserve (DELWP 2019d).
Diamond Firetail	<i>Stagonopleura guttata</i>	There are 69 records of Diamond Firetail from the local area, the most recent in 2011 (DELWP 2019d). Diamond Firetail generally prefers woodland habitats, but is also associated with grassland habitats as well.
Bullant	<i>Myrmecia</i> sp. 17	Although there are only four records listed in the VBA (2019d), the species has large areas of suitable habitat within the study area.

Note. * Those species with the highest likelihood of occurrence.

Recommendation

Targeted surveys have been undertaken for the nationally significant Golden Sun Moth and Striped Legless Lizard to ascertain their presence within the study area, with the results of these surveys provided in a separate report (Ecology and Heritage Partners 2018b). The survey methodology for Striped Legless Lizard is also suitable for the detection of the Fat-tailed Dunnart (and other small mammals).

Based on habitat condition, and the proximity of previous records, there is also low to moderate quality habitat within the study area for the State-significant Speckled Warbler *Chthonicola sagittatus*, Barking Owl

Ninox connivens connivens and Crested Bellbird *Oreoica gutturalis gutturalis* as well as the Regionally significant Fat-tailed Dunnart and Spotted Harrier *Circus assimilis*.

Targeted surveys for State significant fauna would assist to determine their presence within the study area. However, based on existing assessment and approval requirements under the P&E Act, further surveys for these species is not currently required to determine offset implications.

3.4.3 Communities

Two FFG Act-listed ecological communities are present in the study area:

- Western (Basalt) Plains Grassland Community; and,
- Rocky Chenopod Open Scrub Community,

These communities correspond to areas of the Rocky Chenopod Woodland EVC and Plains Grassland EVC mapped in Property 16 within the study area and meets the relevant description and characteristics described for these communities (DELWP 2019h).

3.4.3.1 Western (Basalt) Plains Grassland

One remnant of habitat zone PG4 and all of PG7, PG8 and PG9 meet the description of the State significant *Western (Basalt) Plains Grassland* (WPBG) ecological community (Figure 2). Patches PG8 and PG9 were of the highest quality, were contiguous with each other and other larger remnants of vegetation in Property 16, and supported high native species diversity (Plate 21; Plate 22).



Plate 21. WPBG (PG8) within the study area (Ecology and Heritage Partners Pty Ltd 25/10/2017).



Plate 22. WPBG (PG8) within the study area (Ecology and Heritage Partners Pty Ltd 25/10/2017).

A total of 15.835 hectares of WPBG was recorded within the study area, with 14.456 hectares located in Property 16, and 1.379 hectares situated in Property 11 (Figure 2). All areas of this community are proposed to be retained.

3.4.3.2 *Rocky Chenopod Open Scrub Community*

Habitat zones RCW3, RCW5 and RCW5 meet the description of the State significant Rocky Chenopod Open Scrub Community ecological community (Plate 9; Plate 10) (Figure 2c). A total of 27.799 hectares of the Rocky Chenopod Open Scrub Community ecological community is present. All areas of this community are proposed to be retained.

3.5 Targeted Spiny Rice-flower Survey

A total of 201 Spiny Rice-flower individuals were recorded in Property 11 (Figure 3).

Although the targeted surveys were undertaken during the known flowering period when the species was known to be flowering within the locality, no other specimens were recorded on Properties 4b, 5, 9, 10, or 15 in areas considered to support suitable habitat.

DRAFT

4 LEGISLATIVE AND POLICY IMPLICATIONS

The implications of the project with regards to relevant environmental legislation and policy is outlined below. It should be noted that implications under the EPBC Act, *Planning and Environment Act 1987*, *Wildlife Act 1975*, *Water Act 1989* and *Catchment and Land Protection Act 1994* are detailed in Ecology and Heritage Partners (2018a).

4.1

4.1 *Flora and Fauna Guarantee Act 1988 (Victoria)*

The FFG Act is the primary legislation dealing with biodiversity conservation and sustainable use of native flora and fauna in Victoria. Proponents are required to apply for an FFG Act Permit to 'take' listed and/or protected flora species, listed vegetation communities and listed fish species in areas of public land (i.e. within road reserves, drainage lines and public reserves). An FFG Act permit is generally not required for removal of species or communities on private land, or for the removal of habitat for a listed terrestrial fauna species.

There is suitable habitat within the study area for several 'listed' and 'protected' flora and fauna species under the FFG Act (Appendix 2.1; Section 3.4). Further, two FFG Act communities are present within the study area (Section 3.4.3).

4.1.1 Implications

There is suitable habitat within the study area for several species listed or protected under the FFG Act (Section 3.4). However, the study area is privately owned, as such a permit under the FFG Act is not required, unless impacts to FFG Act listed matters are impacted on public land (i.e. road reserves). No FFG Act listed communities will be impacted by the proposed development. If required, the proponent should allow up to six weeks to obtain a FFG Act permit through DELWP.

4.2 *Environment Effects Act 1978*

The *Environment Effects Act 1978* (EE Act) provides for assessment of proposed actions that are capable of having a significant effect on the environment via the preparation of an Environment Effects Statement (EES). A project with potential adverse environmental effects that, individually or in combination, could be significant in a regional or State context should be referred. An action may be referred for an EES decision where:

- one of the following occurs:
 - Potential clearing of 10 hectares or more of native vegetation from an area that:
 - is of an EVC identified as endangered by DELWP;
 - is, of Very High conservation significance; or,

- is not authorised under an approved Forest Management Plan or Fire Protection Plan.
 - Potential long-term loss of a significant proportion (1-5% depending on conservation status of species) of known remaining habitat or population of a threatened species within Victoria.
- or where two or more of the following occur:
 - Potential clearing of 10 hectares or more of native vegetation, unless authorised under an approved Forest Management Act or Fire Protection Plan;
 - Matters listed under the FFG Act:
 - Potential loss of a significant area of a listed ecological community;
 - Potential loss of a genetically important population of an endangered or threatened species;
 - Potential loss of critical habitat; or,
 - Potential significant effects on habitat values of a wetland supporting migratory birds.

4.2.1 Implications

Table 7 details individual environment effects that may warrant a referral, whilst Table 8 details criteria where a combination of two or more environmental effects may warrant a referral.

Table 7. Individual potential environmental effects that may warrant a referral

Item	Criteria	Response
1	<p>Potential clearing of 10 ha or more of native vegetation from an area that:</p> <ul style="list-style-type: none"> • is of an Ecological Vegetation Class identified as <u>endangered</u> by the DELWP (in accordance with Appendix 2 of Victoria’s Native Vegetation Management Framework); <u>or</u> • is, or is likely to be, of <u>Very High</u> conservation significance (as defined in accordance with Appendix 3 of Victoria’s Native Vegetation Management Framework); and, • is not authorised under an approved Forest Management Plan or Fire Protection Plan. 	<p>A total of 12.069 hectares of native vegetation is to be impacted as part of the development, which consists of 5.3201 hectares of ‘Current Wetland’, and a total of 6.7489 hectares of native vegetation as mapped during the field assessments.</p> <p>The vegetation within the 5.3201 hectares covered by the Current Wetland layer did not meet the thresholds that define a patch of native vegetation despite the field assessments being undertaken during a suitable seasonal timeframe to assess the presence of seasonally dominant wetland vegetation communities. Further, a hydrological assessment of the wetland concluded that a more suitable representation of the area likely to support a shallow ephemeral wetland is approximately 2 hectares in size (Water Technology 2019). An application to reassess the extent of the Current Wetland modelled within Property 5 is currently being prepared for submission to DELWP.</p> <p>The vegetation within the 6.7489 hectares of mapped vegetation consists of:</p> <ul style="list-style-type: none"> • 6.7489 hectares of EVCs classified as Endangered (Plains Grassland); and, • 0.00 hectares of Very High conservation significance vegetation. <p>The total extent of native vegetation (including the Current Wetland layer) proposed to be removed is below the threshold</p>

Item	Criteria	Response
2	Potential long-term loss of a significant proportion (e.g. 1 to 5 percent depending on the conservation status of the species) of known remaining habitat or population of a threatened species within Victoria	<p>that triggers a recommendation for referral under the EE Act.</p> <p>Flora</p> <p>Six State-significant flora listed on the DELWP Advisory List (Spiny Rice-flower, Fragrant Saltbush, Black Roly-poly, Slender Bindweed, Melbourne Yellow-gum and Bacchus Marsh Wattle) were recorded during the site assessment.</p> <p>Fragrant Saltbush was widely recorded throughout the study area, with approximately 400 specimens estimated to occur. The majority of these occur in Property 16 which will be retained as part of the proposed development. As the species is widespread through much of the Victorian Volcanic Plain bioregion within Victoria, the long-term loss will not exceed 1-5% of overall habitat.</p> <p>All Spiny Rice-flower specimens recorded during the targeted surveys, and all specimens of Black Roly-poly, Slender Bindweed and Melbourne Yellow-gum observed are in areas proposed to be retained.</p> <p>Some planted specimens of Bacchus Marsh Wattle may be impacted by the development. However, the long-term loss will not exceed 1-5% of overall habitat as the majority of these specimens are located in areas proposed to be retained.</p> <p>There is potential habitat for a range of State-significant flora within the study area. However, due to the highly modified, agricultural nature of the study area, landscape context and the proximity of previous records, additional State significant flora species listed on DELWP's advisory list are considered unlikely to occur within the study area.</p> <p>Fauna</p> <p>Approximately 22.197 hectares of confirmed habitat for one State-significant fauna listed on the DELWP Advisory List (Golden Sun Moth) is proposed to be impacted, with approximately 28 hectares of confirmed habitat retained.</p> <p>Given the known distribution of Golden Sun Moth within Victoria, the loss of 22.197 hectares does not exceed 1-5% of overall habitat.</p> <p>There is also suitable foraging habitat within the study area for numerous State listed bird species.</p> <p>These species are considered likely to utilise woodland habitat within Properties, 1, 2, 8 and 16 which are proposed to be retained. Further, higher quality foraging habitat is located nearby at Long Forest Flora and Fauna Reserve, and Lerderderg State Park.</p> <p>As the majority of suitable foraging habitat for these, and other bird species will be retained as part of the development, it is not considered that a significant proportion of known remaining habitat will be impacted.</p> <p>Striped Legless Lizard was recorded within areas of potential habitat during targeted surveys, and based on the results of the targeted survey, there is considered a low likelihood that an extant population of the species occurs within the study area.</p> <p>As such, it is considered that there will not be a long-term loss of known remaining habitat for any threatened fauna species that will exceed 1-5% of overall habitat.</p>
3	Potential long-term change to the ecological character of a wetland listed under the	The nearest wetland listed under the Ramsar Convention is located 35 km south of the site (Port Phillip Bay – western Shoreline and

Item	Criteria	Response
	Ramsar Convention or in 'A Directory of Important Wetlands in Australia'	Bellarine Peninsula). The nearest wetland in the 'Directory of Important Wetlands' is the Lerderderg River, located approximately 3.3 km west of the study area. Provided management practices and construction techniques are consistent with Construction Techniques for Sediment Pollution Control (EPA 1991) and Environmental Guidelines for Major Construction Sites (EPA 1996), the proposed action is unlikely to impact the ecological character of any Ramsar site or 'important wetland'.
4	Potential extensive or major effects on the health or biodiversity of aquatic, estuarine or marine ecosystems, over the long term	There are no estuarine, marine or permanent aquatic ecosystems within the study area. As such, there is not considered to be any extensive or major effects to any aquatic, estuary or marine system over the long-term.

Table 8. A combination of two or more environmental effects that may warrant a referral

Item	Criteria	Response
1	Potential clearing of 10 hectares or more of native vegetation, unless authorised under an approved Forest Management Plan or Fire Protection Plan	A total of 12.069 hectares of native vegetation is to be impacted as part of the development, which consists of 5.3201 hectares of 'Current Wetland', and a total of 6.7489 hectares of native vegetation as mapped during the field assessments.
2	Matters listed under the <i>Flora and Fauna Guarantee Act 1988</i> : <ul style="list-style-type: none"> potential loss of a significant area of a listed ecological community; or, potential loss of a genetically important population of an endangered or threatened species (listed or nominated for listing), including as a result of loss or fragmentation of habitats; or potential loss of critical habitat; or potential significant effects on habitat values of a wetland supporting migratory bird species 	There are no FFG Act-listed communities being impacted within the study area. It is not considered that the study area supports a genetically important population of, or critical habitat for any FFG Act-list species. Migratory and wetland bird species are not considered likely to utilise ephemeral wetlands within the study area on a regular or permanent basis given higher quality foraging habitat is located nearby at Merrimu Reservoir. Based on the above, we do not consider that any of the thresholds relating to FFG Act criteria have been exceeded.
3	Potential extensive or major effects on landscape values of regional importance, especially where recognised by a planning scheme overlay or within or adjoining land reserved under the <i>National Parks Act 1975</i>	Long Forest Nature Conservation Reserve is covered by an ESO3 overlay. No impacts to the Long Forest Reserve are proposed. An SLO1 affects a portion of land south of Buckleys Road that intersects with an area of retained native vegetation within Property 8. As such, it is not considered that the proposed development will result in extensive or major effects to landscapes of regional importance. The proposal is also not within or adjoining land reserved under the <i>National Parks Act 1975</i> . Based on the above, we do not consider this threshold to be exceeded.

Based on a review of ecological impacts associated with the proposed development, it is Ecology and Heritage Partners' position that a referral under the EES Act is not required based on ecological impacts alone, as:

- None of the thresholds relating to any of the ecological criteria identified in Table 7 have been exceeded; and,
- None of the thresholds relating to any of the ecological criteria identified in Table 8 have been exceeded.

It should be noted that Ecology and Heritage Partners' have not undertaken a detailed assessment of other non-ecological referral criteria detailed in DSE (2006).

DRAFT

5 MITIGATION MEASURES

For the removal of vegetation that falls under all assessment pathways, the Guidelines (DELWP 2017a) require the responsible authority to consider whether the applicant has demonstrated avoidance and minimisation of impacts to native vegetation.

5.1 Avoidance and Minimisation Statement

The proposed development plan will impact upon a large proportion of the study site, with the majority or areas proposed to be impacted comprised of cropped agricultural land devoid of native vegetation. Of the 72.791 hectares of native vegetation mapped within Properties 1-16, impacts to a total of 60.443 hectares of the highest quality native vegetation will be avoided. Of the 12.069 hectares of native vegetation proposed to be impacted, 5.323 hectares comprises a modelled wetland that is highly modified, has been subjected to multiple cropping events in recent years, and is considered highly unlikely to continue to support seasonal wetland vegetation due to the alterations to the historical hydrological influences caused by agricultural practices (ploughing, tilling cropping).

All of the vegetation on Property 16 (42.894 hectares) is proposed for use as an offset site, and will be protected and managed in order to enhance to existing biodiversity values that occur within the site and surrounds. All scattered trees and large old trees within patches that are located within the development footprint (properties 1-15) will be retained. In addition, the population of Spiny Rice-flower present on Property 11 will be retained.

In the context of the contribution to Victoria's biodiversity that the native vegetation makes, the above avoidance and minimisation measures are considered appropriate in the context of the project, and broader ecological values within and adjacent to the study area.

5.2 Best Practice Mitigation Measures

Recommended measures to mitigate impacts upon terrestrial and aquatic values present within the study area may include:

- Minimise impacts to native vegetation and habitats through construction and micro-siting techniques, including fencing retained areas of native vegetation. If indeed necessary, trees should be lopped or trimmed rather than removed. Similarly, soil disturbance and sedimentation within wetlands should be avoided or kept to a minimum, to avoid, or minimise impacts to fauna habitats;
- All contractors should be aware of ecologically sensitive areas to minimise the likelihood of inadvertent disturbance to areas marked for retention. Habitat Zones (areas of sensitivity) should be included as a mapping overlay on any construction plans;
- Tree Retention Zones (TRZs) should be implemented to prevent indirect losses of native vegetation during construction activities (DSE 2011). A TRZ applies to a tree and is a specific area above and below the ground, with a radius 12 x the DBH. At a minimum standard a TRZ should consider the following:
 - A TRZ of trees should be a radius no less than two metres or greater than 15 metres;

- Construction, related activities and encroachment (i.e. earthworks such as trenching that disturb the root zone) should be excluded from the TRZ;
 - Where encroachment exceeds 10% of the total area of the TRZ, the tree should be considered as lost and offset accordingly;
 - Directional drilling may be used for works within the TRZ without being considered encroachment. The directional bore should be at least 600 millimetres deep;
 - The above guidelines may be varied if a qualified arborist confirms the works will not significantly damage the tree (including stags / dead trees). In this case the tree would be retained and no offset would be required; and,
 - Where the minimum standard for a TRZ has not been met an offset may be required.
- Consideration of Water Sensitive Urban Design techniques such as stormwater treatment wetlands, bio-retention systems, porous paving or swales;
 - Where possible, construction stockpiles, machinery, roads, and other infrastructure should be placed away from areas supporting native vegetation, LOTs and/or wetlands;
 - Ensure that best practice sedimentation and pollution control measures are undertaken at all times, in accordance with Environment Protection Authority guidelines (EPA 1991; EPA 1996; Victorian Stormwater Committee 1999) to prevent offsite impacts to waterways and wetlands; and,
 - As indigenous flora provides valuable habitat for indigenous fauna, it is recommended that any landscape plantings that are undertaken as part of the proposed works are conducted using indigenous species sourced from a local provenance, rather than exotic deciduous trees and shrubs.

In addition to these measures, the following documents should be prepared and implemented prior to any construction activities:

- Construction Environmental Management Plan (CEMP). The CEMP should include specific species/vegetation conservation strategies, daily monitoring, sedimentation management, and site specific rehabilitation plans. A weed management plan is likely to be required as a component of the CEMP.

5.3 Offset Impacts

Ecology and Heritage Partners are a DELWP accredited OTC offset broker and BushBroker site assessor.

Based on an assessment of the native vegetation to be retained and protected in Property 16, and the gain that can be generated through the permanent protection of native vegetation via an on-title security mechanism (i.e. Section 69 agreement under the *Conservation, Forests and Lands Act 1987*), and ongoing management and enhancement activities, the majority of the offset obligation can be met on-site (Appendix 4.2).

The following offset credits relevant to the development can be generated via the management and security of 42.894 hectares of native vegetation (Appendix 4.2):

- 12.973 General HUs; and,
- 12.114 species habitat units for Bacchus Marsh Wattle;

- 4.757 species habitat units for Small Golden Moths;
- 12.109 species habitat units for Fragrant Saltbush,
- 12.109 species habitat units for Heath Spear-grass; and,
- 12.109 species habitat units for Melbourne Yellow-gum.

This leaves a deficit of:

- 9.675 species units of habitat for Basalt Podolepis; and,
- 4.983 species units of habitat for Small Golden Moths.

Ecology and Heritage Partners are currently preparing a separate offset strategy to demonstrate how these obligations can and will be secured.

DRAFT

6 FURTHER REQUIREMENTS

Further requirements associated with development of the study area, as well as additional studies or reporting that may be required, are provided below (Table 8).

Table 8. Further requirements associated with development of the study area.


Relevant Legislation	Implications	Further Action
<i>Flora and Fauna Guarantee Act 1988</i>	There is suitable habitat within the study area for several species listed or protected under the FFG Act (Section 3.4). However, the study area is privately owned, as such a permit under the FFG Act is not required, unless impacts to FFG Act listed matters are impacted on public land (i.e. road reserves). No FFG Act listed communities will be impacted by the proposed development. If required, the proponent should allow up to six weeks to obtain a FFG Act permit through DELWP.	No further action required.
<i>Environment Effects Act 1978</i>	<p>A total of 12.069 hectares of native vegetation is to be impacted as part of the development, which consists of 5.3201 hectares of 'Current Wetland', and a total of 6.7489 hectares of native vegetation as mapped during the field assessments.</p> <p>A hydrological assessment of the wetland concluded that a more suitable representation of the area likely to support a shallow ephemeral wetland is approximately 2 hectares in size (Water Technology 2019).</p> <p>It is considered that there will not be a long-term loss of known remaining habitat for any threatened fauna species that will exceed 1-5% of overall habitat.</p>	Based on a review of ecological impacts associated with the proposed development, it is Ecology and Heritage Partners' position that a referral under the EES Act is not required based on ecological impacts alone.

REFERENCES

- ALA 2017. Atlas of Living Australia. [www Document]. URL: <http://www.ala.org.au/>. NCRIS – National Research Infrastructure for Australia. Commonwealth of Australia.
- BL&A 2010. Bences Road, Bacchus marsh. Habitat Hectare Assessment. Report prepared for ABC Project Management Pty Ltd. Brett Lane & Associates Pty Ltd. January 2010.
- BL&A 2012. 289 Bences Road, Bacchus Marsh: Offset Options. Report prepared for Queensland Investment Corporation. Brett Lane & Associates Pty Ltd July 2012.
- Christidis, L. and Boles, W.E 2008. Systematics and Taxonomy of Australian Birds. CSIRO Publishing, Collingwood, Victoria.
- Cogger, H. G (Ed). 1996. Reptiles and Amphibians of Australia. 5th Edition. Reed Books Australia, Victoria.
- Cogger, H. G., Cameron, E. E. And Cogger, H. M. 1983. *Volume 1 of Zoological Catalogue of Australia: Amphibia and Reptilia*. Australian Government Publishing Service, Canberra, ACT.
- Cogger, H.G., Cameron, E.E., Sadler, R.A. and Egger P., 1993. The Action Plan for Australian Reptiles. Australian Nature conservation Agency, Canberra, ACT.
- DELWP 2016. *Flora and Fauna Guarantee Act 1988* Protected Flora List – December 2016. Victorian Department of Environment, Land, Water and Planning. Melbourne, Victoria.
- DELWP 2017a. Guidelines for the removal, destruction or lopping of native vegetation. The State of Victoria Department of Environment, Land, Water and Planning . Melbourne, December 2017.
- DELWP 2019a. Native Vegetation Information Management Tool [www Document]. URL: <https://nvim.delwp.vic.gov.au/>. Victorian Department of Environment, Land, Water and Planning, Melbourne, Victoria.
- DELWP 2019b. NatureKit Map [www Document]. URL: <http://maps.biodiversity.vic.gov.au/viewer/?viewer=NatureKit>. Victorian Department of Environment, Land, Water and Planning, Melbourne, Victoria.
- DELWP 2019c. Ecological Vegetation Class (EVC) Benchmarks for each Bioregion [www Document]. URL: <http://www.depi.vic.gov.au/environment-and-wildlife/biodiversity/evc-benchmarks#bioregionname>. Victorian Department of Environment, Land, Water and Planning, Melbourne, Victoria.
- DELWP 2019d. Victorian Biodiversity Atlas. Sourced from GIS layers: “VBA_FLORA25”, “VBA_FLORA100”, “VBA_FAUNA25”, “VBA_FAUNA100”, July 2018. Victorian Department of Environment, Land, Water and Planning, Melbourne, Victoria.
- DELWP 2018. *Flora and Fauna Guarantee Act 1988* Threatened List – March 2017. Victorian Department of Environment, Land, Water and Planning, Melbourne, Victoria.
- DELWP 2019f. Planning Maps Online [www Document]. URL: <http://services.land.vic.gov.au/maps/pmo.jsp>. Victorian Department of Environment, Land, Water and Planning, Melbourne, Victoria.

- DELWP 2019g. Planning Schemes Online [www Document]. URL: <http://planningschemes.dpcd.vic.gov.au>. Victorian Department of Environment, Land, Water and Planning, Melbourne, Victoria.
- DELWP 2019h. *Flora and Fauna Guarantee Act 1988* – Threatened List: Characteristics of Threatened Communities [WWW Document]. URL: < <https://www.environment.vic.gov.au/conserving-threatened-species/flora-and-fauna-guarantee-act-1988> >.
- DELWP 2017b. Assessor’s handbook. Applications to remove, destroy or lop native vegetation. Version 1.0. The State of Victoria Department of Environment, Land, Water and Planning . Melbourne, December 2017.
- DEPI 2014. Advisory List of Rare or Threatened Plants in Victoria. Victorian Department of Environment and Primary Industries, Melbourne, Victoria.
- DEWHA 2009. Significant impact guidelines for the critically endangered Spiny Rice-flower (*Pimelea spinescens* subsp. *spinescens*). Nationally threatened species and ecological communities EPBC Act Policy Statement 3.11. Commonwealth of Australia, 2009.
- DoE 2013. Significant Impact Guidelines 1.1. Matters of National Environmental Significance. Commonwealth Department of the Environment, Canberra, ACT.
- DoEE 2018. Protected Matters Search Tool: Interactive Map [www Document]. URL: <<http://www.environment.gov.au/epbc/pmst/>>. Commonwealth Department of the Environment and Energy, Canberra, ACT.
- DSE 2004. Vegetation quality assessment manual: Guidelines for applying the habitat hectares scoring method. Version 1.3. Victorian Department of Sustainability and Environment, Melbourne Victoria
- DSE 2006. Ministerial guidelines for assessment of environmental effects under the *Environment Effects Act 1978*. Seventh Edition, 2006. Published by the Victorian Government Department of Sustainability and Environment, Melbourne. June 2006
- DSE 2009. Advisory list of Threatened Invertebrate Fauna in Victoria – 2009. Victorian Department of Sustainability and Environment, Melbourne, Victoria.
- DSE 2010a. Biodiversity Precinct Structure Planning Kit. Published by the Victorian Government Department of Sustainability and Environment. Melbourne. May 2010.
- DSE 2010b. National Recovery Plan for the Small Golden Moths Orchid *Diuris basaltica*. Published by the Victorian Government Department of Sustainability and Environment, Melbourne. October 2010.
- DSE 2010c. National Recovery Plan for the Basalt Peppercross *Lepidium hyssopifolium*. Published by the Victorian Government Department of Sustainability and Environment, Melbourne. July 2010.
- DSE 2013. Advisory List of Threatened Vertebrate Fauna in Victoria. Victorian Department of Sustainability and Environment, Melbourne, Victoria.
- Duncan, A., Baker, G.B. and Montgomery, N. (Eds) 1999. The Action Plan for Australian Bats. Environment Australia. Canberra, ACT.
- Ecology and Heritage Partners Pty Ltd 2013a. Flora and Fauna Assessment, and Net Gain Assessment at Long Forest Estate, Merrimu, Victoria. Report prepared for QOD Property Group. December 2013.

- Ecology and Heritage Partners Pty Ltd 2013b. Targeted Flora and Fauna Surveys at Long Forest Estate, Merrimu, Victoria. Report prepared for QOD Property Group. December 2013.
- Ecology and Heritage Partners Pty Ltd 2018a. Ecological Assessment: Bacchus Marsh Development Project, Victoria. Report prepared for Bacchus Marsh Developments Pty Ltd. July 2018.
- Ecology and Heritage Partners Pty Ltd 2018b. Targeted Surveys for Golden Sun Moth *Synemon plana* and Striped Legless Lizard *Delma impar*: Bacchus Marsh Development Project, Bacchus Marsh, Victoria. Report prepared for Bacchus Marsh Developments Pty Ltd. June 2018.
- EPA 1991. Construction Techniques for Sediment Pollution Control. Published document prepared by the Victorian Environment Protection Authority, Melbourne, Victoria.
- EPA 1996. Environmental Guidelines for Major Construction Sites. Published document prepared by the Victorian Environmental Protection Authority, Melbourne, Victoria.
- Garnett, S., J. Szabo and G. Dutson 2011. The Action Plan for Australian Birds 2010. CSIRO Publishing, Collingwood, Victoria.
- Gullan, P 2017f. Illustrated Flora Information System of Victoria (IFLISV). Viridans Pty Ltd, Victoria.
- Menkhorst, P. and Knight, F. 2004. A Field Guide to the Mammals of Australia . 2nd Edition. Oxford University Press, Victoria.
- Moorabool Shire 2017. Amendment C81 Bacchus Marsh Urban Growth Framework. [www Document]. URL: <https://www.moorabool.vic.gov.au/consultations/amendment-c81-bacchus-marsh-urban-growth-framework>. Moorabool Shire Council, Victoria.
- Nelson, J. S. 1994. Fishes of the World, 3rd Edition. John Wiley & Sons, New York, USA.
- Paul Kelly & Associates 2014. Site Assessments – Bacchus Marsh. Report prepared for ABC Project Management. July 2014.
- Sands, D.P.A. and New, T.R. 2002. The Action Plan for Australian Butterflies, Environment Australia, Canberra, ACT.
- SEWPaC 2012. *Environment Protection and Biodiversity Conservation Act 1999* Environmental Offsets Policy. Commonwealth Department of Environment, Water, Population and Communities, Canberra, ACT.
- Strahan, R. (Ed) 1995. The Mammals of Australia. Reed Books, Sydney, NSW.
- Tyler, M.J. 1997. The Action Plan for Australian Frogs. Wildlife Australia: Canberra, ACT.
- Victorian Urban Stormwater Committee 1999. Urban Stormwater: Best Practice Environmental Management Guidelines. CSIRO, Collingwood, Victoria.
- Viridans (2014). Victorian Fauna Database. Viridans Biological Databases, Bentleigh East, Victoria.
- Water Technology 2019. Memorandum: Merrimu Water Balance. Prepared for Stamoulis Property Group. 23 January 2019.
- Woinarski J. C. Z., Burbidge A. A. and Harrison P. 2014. The action plan for Australian mammals 2012. CSIRO Publishing, Collingwood, Victoria.



Legend

- Study Area
- Railway
- Freeway
- Major Road
- Collector Road
- Minor Road
- Minor Watercourse
- Major Watercourse
- Permanent Waterbody
- Land Subject to Inundation
- Wetland/Swamp
- Parks and Reserves
- Crown Land
- Localities





Figure 1
Location of the study area
Ecological Assessments for the Bacchus Marsh Development Project



VicMap Data: The State of Victoria does not warrant the accuracy or completeness of information in this publication and any person using or relying upon such information does so on the basis that the State of Victoria shall bear no responsibility or liability whatsoever for any errors, faults, defects or omissions in the information.




Figure 2 Overview Legend

Ecological features
Ecological Assessments for the Bacchus Marsh Development Project

- Study Area
- Proposed lot division
- Golden Sun Moth Habitat
- ▲ Lot in Patch retained
- Scattered Tree retained

VROTs


- Melbourne Yellow-Gum
- Slender Bindweed
- Fragrant Saltbush
- Black Roly-Poly
- Spiny Rice-flower

Ecological Vegetation Classes

- Grassy Woodland EVC 175
- Plains Grassland EVC 132
- Plains Grassy Wetland EVC 125
- Rocky Chenopod Woodland EVC 64
- Current Wetland
- Retained vegetation
- Removed vegetation

FFG Act listed community

- Western (Basalt) Plains Grassland
- Rocky Chenopod Open Scrub



VicMap Data: The State of Victoria does not warrant the accuracy or completeness of information in this publication and any person using or relying upon such information does so on the basis that the State of Victoria shall bear no responsibility or liability whatsoever for any errors, faults, defects or omissions in the information.





Figure 2a
Ecological features
Ecological Assessments for the Bacchus Marsh Development Project

Legend

- Study Area
- Golden Sun Moth Habitat
- ✪ Scattered Tree retained
- VROTs**
- Fragrant Saltbush

- Ecological Vegetation Classes**
- Grassy Woodland EVC 175
 - Plains Grassland EVC 132
 - Plains Grassy Wetland EVC 125
 - Rocky Chenopod Woodland EVC 64
 - Current Wetland
 - Retained vegetation
 - Removed vegetation







Figure 2b
Ecological features
Ecological Assessments for the Bacchus Marsh Development Project

Legend

- Study Area
- Golden Sun Moth Habitat
- ▲ Lot in Patch retained
- Scattered Tree retained
- VROTs**
- Melbourne Yellow-Gum
- Fragrant Saltbush
- Spiny Rice-flower

- Ecological Vegetation Classes**
- Grassy Woodland EVC 175
 - Plains Grassland EVC 132
 - Retained vegetation
 - Removed vegetation

- FFG Act listed community**
- Western (Basalt) Plains Grassland



VicMap Data: The State of Victoria does not warrant the accuracy or completeness of information in this publication and any person using or relying upon such information does so on the basis that the State of Victoria shall bear no responsibility or liability whatsoever for any errors, faults, defects or omissions in the information.





Figure 2c
Ecological features
Ecological Assessments for the Bacchus Marsh Development Project

Legend

- Study Area
- Proposed lot division
- Golden Sun Moth Habitat
- VROTs**
- Melbourne Yellow-Gum
- Slender Bindweed
- Fragrant Saltbush
- Black Roly-Poly

- Ecological Vegetation Classes**
- Plains Grassland EVC 132
 - Rocky Chenopod Woodland EVC 64
 - Retained vegetation
 - Removed vegetation

- FFG Act listed community**
- Western (Basalt) Plains Grassland
 - Rocky Chenopod Open Scrub



VicMap Data: The State of Victoria does not warrant the accuracy or completeness of information in this publication and any person using or relying upon such information does so on the basis that the State of Victoria shall bear no responsibility or liability whatsoever for any errors, faults, defects or omissions in the information.





Figure 3
Spiny Rice-flower
Survey Results
Ecological Assessments
for the Bacchus Marsh
Development Project

Legend

- Study Area
- Spiny Rice-flower
- Ecological Vegetation Classes**
- Plains Grassland EVC 132
- FFG Act listed community**
- Western (Basalt) Plains Grassland



VicMap Data: The State of Victoria does not warrant the accuracy or completeness of information in this publication and any person using or relying upon such information does so on the basis that the State of Victoria shall bear no responsibility or liability whatsoever for any errors, faults, defects or omissions in the information.




- Legend**
- Study Area
- Significant flora**
- Arching Flax-lily
 - Austral Tobacco
 - Bacchus Marsh Wattle
 - Black Roly-poly
 - Black-tip Greenhood
 - Branching Groundsel
 - Brittle Greenhood
 - Buloke
 - Cane Spear-grass
 - Curved Rice-flower
 - Forked Rice-flower
 - Fragrant Saltbush
 - Heath Spear-grass
 - Leafless Bluebush
 - Melbourne Yellow-gum
 - Narrow-leaf Wax-flower
 - ▲ Rough Wattle
 - ▲ Rye Beetle-grass
 - ▲ Slender Bindweed
 - ▲ Small Golden Moths
 - ▲ Smooth Nardoo
 - ▲ Snowy Mint-bush
 - ▲ Spiny Rice-flower
 - ▲ Werribee Blue-box
 - + Western Golden-tip

Figure 4
 Previously documented significant flora within 5km of the study area
Ecological Assessments for the Bacchus Marsh Development Project



VBA 2017. Victorian Biodiversity Atlas. // Sourced from: 'VBA_FLORA25' and 'VBA_FLORA100', February 2017 © The State of Victoria, Department of Environment, Land, Water and Planning. Records prior to 1949 not shown.
 VicMap Data: The State of Victoria does not warrant the accuracy or completeness of information in this publication and any person using or relying upon such information does so on the basis that the State of Victoria shall bear no responsibility or liability whatsoever for any errors, faults, defects or omissions in the information.
 10937_Fig04_SigFlora_f9794 31/01/2019 psorensen



- Legend**
- Study Area
- Significant fauna**
- Australasian Shoveler
 - Azure Kingfisher
 - Barking Owl
 - Black Falcon
 - Black-eared Cuckoo
 - Blue-billed Duck
 - Brown Toadlet
 - Brown Treecreeper (south-eastern ssp.)
 - Brush-tailed Phascogale
 - Bullant
 - Caddisfly
 - Caspian Tern
 - Chestnut-rumped Heathwren
 - Common Dunnart
 - Crested Bellbird
 - Diamond Firetail
 - ▲ Eastern Great Egret
 - ▲ Eastern Pygmy-possum
 - ▲ Eastern Snake-necked Turtle
 - ▲ Freckled Duck
 - ▲ Glossy Ibis
 - ▲ Golden Sun Moth
 - ▲ Grey-headed Flying-fox
 - ▲ Growling Grass Frog
 - + Gull-billed Tern
 - + Hardhead
 - + Hooded Robin
 - + Lace Monitor
 - + Latham's Snipe
 - + Musk Duck
 - + Nankeen Night Heron
 - + Pied Cormorant
 - ⌘ Powerful Owl
 - ⌘ Royal Spoonbill
 - ⌘ Speckled Warbler
 - ⌘ Spotted Harrier
 - ⌘ Swift Parrot
 - ⌘ White-bellied Sea-Eagle
 - ⌘ White-throated Needletail

Figure 5
 Previously documented significant fauna within 5km of the study area
Ecological Assessments for the Bacchus Marsh Development Project



APPENDICES

DRAFT

APPENDIX 1

Appendix 1.1 – Rare or Threatened Categories for Listed Victorian Taxa

Table A1.1. Rare or Threatened categories for listed Victorian taxa.

Rare or Threatened Categories
Conservation Status in Australia (Based on the EPBC Act 1999)
EX - Extinct: Extinct is when there is no reasonable doubt that the last individual of the species has died.
CR - Critically Endangered: A species is critically endangered when it is facing an extremely high risk of extinction in the wild in the immediate future.
EN - Endangered: A species is endangered when it is not critically endangered but is facing a very high risk of extinction in the wild in the near future.
VU - Vulnerable: A species is vulnerable when it is not critically endangered or endangered but is facing a high risk of extinction in the wild in the medium-term future.
R* - Rare: A species is rare but overall is not currently considered critically endangered, endangered or vulnerable.
K* - Poorly Known: A species is suspected, but not definitely known, to belong to any of the categories extinct, critically endangered, endangered, vulnerable or rare.
Conservation Status in Victoria (Based on DEPI 2014, DSE 2009 or 2013)
x - Presumed Extinct in Victoria: not recorded from Victoria during the past 50 years despite field searches specifically for the plant, or, alternatively, intensive field searches (since 1950) at all previously known sites have failed to record the plant.
e - Endangered in Victoria: at risk of disappearing from the wild state if present land use and other causal factors continue to operate.
v - Vulnerable in Victoria: not presently endangered but likely to become so soon due to continued depletion; occurring mainly on sites likely to experience changes in land-use which would threaten the survival of the plant in the wild; or, taxa whose total population is so small that the likelihood of recovery from disturbance, including localised natural events such as drought, fire or landslip, is doubtful.
r - Rare in Victoria: rare but not considered otherwise threatened - there are relatively few known populations or the taxon is restricted to a relatively small area.
k - Poorly Known in Victoria: poorly known and suspected, but not definitely known, to belong to one of the above categories (x, e, v or r) within Victoria. At present, accurate distribution information is inadequate.

Appendix 1.2 – Defining Ecological Significance

Table A1.2. Criteria for defining Ecological Significance ratings for significant flora, fauna and communities.

National Significance
<p>Flora: National conservation status is based on the EPBC Act list of taxa considered threatened in Australia (i.e. extinct, critically endangered, endangered, vulnerable).</p>
<p>Fauna: National conservation status is based on the EPBC Act list of taxa considered threatened in Australia (i.e. Extinct, Critically Endangered, Endangered, Vulnerable). Fauna listed as Extinct, Critically Endangered, Endangered, Vulnerable, or Rare under National Action Plans for terrestrial taxon prepared for DoE: mammals (Woinarski <i>et al.</i> 2014), bats (Duncan <i>et al.</i> 1999), birds (Garnett <i>et al.</i> 2011), reptiles (Cogger <i>et al.</i> 1993), amphibians (Tyler 1997) and butterflies (Sands and New 2002).</p>
<p>Communities: Vegetation communities considered critically endangered, endangered or vulnerable under the EPBC Act and considering vegetation condition.</p>
State Significance
<p>Flora: Threatened taxa listed under the provisions of the FFG Act. Flora listed in the State Government’s Advisory List of Rare or Threatened Plants in Victoria (DEPI 2014).</p>
<p>Fauna: Threatened taxon listed under Schedule 2 of the FFG Act. Fauna listed as Extinct, Critically Endangered, Endangered and Vulnerable on the State Government’s Advisory List of Threatened Vertebrate Fauna in Victoria (DSE 2013). Listed as Lower Risk (Near Threatened, Conservation Dependent or Least concern) or Data Deficient under National Action Plans for terrestrial species prepared for the DoE: mammals (Woinarski <i>et al.</i> 2014), bats (Duncan <i>et al.</i> 1999), birds (Garnett <i>et al.</i> 2011), reptiles (Cogger <i>et al.</i> 1993), amphibians (Tyler 1997) and butterflies (Sands and New 2002).</p>
<p>Communities: Ecological communities listed as threatened under the FFG Act (DELWP 2019g). EVC listed as threatened (i.e. endangered, vulnerable) or rare in a Native Vegetation Plan for a particular bioregion and considering vegetation condition.</p>
Regional Significance
<p>Fauna: Fauna with a disjunct distribution, or a small number of documented recorded or naturally rare in the particular Bioregion in which the study area is located. A particular taxon that is has an unusual ecological or biogeographical occurrence or listed as Lower Risk – Near Threatened, Data Deficient or Insufficiently Known on the State Government’s Advisory List of Threatened Vertebrate Fauna in Victoria (DSE 2013).</p>
<p>Communities: EVC listed as depleted or least concern in a Native Vegetation Plan for a particular bioregion) and considering vegetation condition. EVC considered rare by the author for a particular bioregion.</p>
Local Significance
<p>Local significance is defined as flora, fauna and ecological communities indigenous to a particular area, which are not considered rare or threatened on a national, state or regional level.</p>

Appendix 1.3 – Defining Site Significance

Table A1.3. Criteria for defining Site Significance ratings.

National Significance
<p>A site is of National significance if:</p> <ul style="list-style-type: none"> It regularly supports, or has a high probability of regularly supporting individuals of a taxon listed as ‘Critically Endangered’ or ‘Endangered’ under the EPBC Act and/or under National Action Plans for terrestrial taxon prepared for the DoE. It regularly supports, or has a high probability of supporting, an ‘important population’ as defined under the EPBC Act of one or more nationally ‘vulnerable’ flora and fauna taxon. It is known to support, or has a high probability of supporting taxon listed as ‘Vulnerable’ under National Action Plans. It is known to regularly support a large proportion (i.e. greater than 1%) of a population of a taxon listed as ‘Conservation Dependent’ under the EPBC Act and/or listed as Rare or Lower Risk (near threatened, conservation dependent or least concern) under National Action Plans. It contains an area, or part thereof designated as ‘critical habitat’ under the EPBC Act, or if the site is listed under the Register of National Estate compiled by the Australian Heritage Commission. It is a site which forms part of, or is connected to a larger area(s) of remnant native vegetation or habitat of national conservation significance such as most National Park, and/or a Ramsar Wetland(s).
State Significance
<p>A site is of State significance if:</p> <ul style="list-style-type: none"> It occasionally (i.e. every 1 to 5 years) supports, or has suitable habitat to support taxon listed as ‘Critically Endangered’ or ‘Endangered’ under the EPBC Act and/or under National Action Plans. <ul style="list-style-type: none"> It regularly supports, or has a high probability of regularly supporting (i.e. high habitat quality) taxon listed as ‘Vulnerable’, ‘Near threatened’, ‘Data Deficient’ or ‘Insufficiently Known’ in Victoria (,DSE 2013; DEPI 2014), or species listed as ‘Data Deficient’ or ‘Insufficiently Known’ under National Action Plans. It contains an area, or part thereof designated as ‘critical habitat’ under the FFG Act. It supports, or likely to support a high proportion of any Victorian flora and fauna taxa. It contains high quality, intact vegetation/habitat supporting a high species richness and diversity in a particular bioregion. It is a site which forms part of, or connected to a larger area(s) of remnant native vegetation or habitat of state conservation significance such as most State Parks and/or Flora and Fauna Reserves.
Regional Significance
<p>A site is of Regional significance if:</p> <ul style="list-style-type: none"> It regularly supports, or has a high probability of regularly supporting regionally significant fauna as defined in Table 1.2. Is contains a large population (i.e. greater than 1% or 5%) of flora considered rare in any regional Native Vegetation Plan for a particular bioregion. It supports a fauna population with a disjunct distribution, or a particular taxon that has an unusual ecological or biogeographical occurrence. It is a site which forms part of, or is connected to a larger area(s) of remnant native vegetation or habitat of regional conservation significance such as most Regional Parks and/or Flora and Fauna Reserves.
Local Significance
<p>Most sites are considered to be of at least local significant for conservation, and in general a site of local significance can be defined as:</p> <ul style="list-style-type: none"> An area which supports indigenous flora species and/or a remnant EVC, and habitats used by locally significant fauna species. An area which currently acts, or has the potential to act as a wildlife corridor linking other areas of higher conservation significance and facilitating fauna movement throughout the landscape.

Appendix 1.4 – Vegetation Condition and Habitat Quality

Table A1.4.1 Defining Vegetation Condition ratings.

Criteria for defining Vegetation Condition
<p>High Quality: Vegetation dominated by a diversity of indigenous species, with defined structures (where appropriate), such as canopy layer, shrub layer, and ground cover, with little or few introduced species present.</p>
<p>Moderate Quality: Vegetation dominated by a diversity of indigenous species, but is lacking some structures, such as canopy layer, shrub layer or ground cover, and/or there is a greater level of introduced flora species present.</p>
<p>Low Quality: Vegetation dominated by introduced species, but supports low levels of indigenous species present, in the canopy, shrub layer or ground cover.</p>

Table A1.4.2 Defining Habitat Quality.

Criteria for defining Habitat Quality
<p>High Quality:</p> <ul style="list-style-type: none"> • High degree of intactness (i.e. floristically and structurally diverse), containing several important habitat features such as ground debris (logs, rocks, vegetation), mature hollow-bearing trees, and a dense understorey component. • High species richness and diversity (i.e. represented by a large number of species from a range of fauna groups). • High level of foraging and breeding activity, with the site regularly used by native fauna for refuge and cover. • Habitat that has experienced, or is experiencing low levels of disturbance and/or threatening processes (i.e. weed invasion, introduced animals, soil erosion, salinity). • High contribution to a wildlife corridor, and/or connected to a larger area(s) of high quality habitat. • Provides known, or likely habitat for one or more rare or threatened species listed under the EPBC Act, FFG Act, or species considered rare or threatened according to DEPI 2014; DSE 2009 or 2013.
<p>Moderate Quality:</p> <ul style="list-style-type: none"> • Moderate degree of intactness, containing one or more important habitat features such as ground debris (logs, rocks, vegetation), mature hollow-bearing trees, and a dense understorey component. • Moderate species richness and diversity - represented by a moderate number of species from a range of fauna groups. • Moderate levels of foraging and breeding activity, with the site used by native fauna for refuge and cover. • Habitat that has experienced, or is experiencing moderate levels of disturbance and/or threatening processes. • Moderate contribution to a wildlife corridor, or is connected to area(s) of moderate quality habitat. • Provides potential habitat for a small number of threatened species listed under the EPBC Act, FFG Act, or species considered rare or threatened according to DEPI 2014; DSE 2009 or 2013.
<p>Low Quality:</p> <ul style="list-style-type: none"> • Low degree of intactness, containing few important habitat features such as ground debris (logs, rocks, vegetation), mature hollow-bearing trees, and a dense understorey component. • Low species richness and diversity (i.e. represented by a small number of species from a range of fauna groups). • Low levels of foraging and breeding activity, with the site used by native fauna for refuge and cover. • Habitat that has experienced, or is experiencing high levels of disturbance and/or threatening processes. • Unlikely to form part of a wildlife corridor, and is not connected to another area(s) of habitat. • Unlikely to provide habitat for rare or threatened species listed under the EPBC Act, FFG Act, or considered rare or threatened according to DEPI 2014; DSE 2009 or 2013.

Appendix 1.5 – Offsets and Exemptions

Table A1.5.1. Calculation of Biodiversity Equivalence Scores and General or Specific Offsets (DEPI 2013)

Pathway	Biodiversity Assessment Tools	Information Source
Low Risk-based pathway	Condition Score	Modelled data, NVIM Tool (DELWP 2017a)
	Habitat Hectares	= Condition Score x Extent (ha)
	Strategic Biodiversity Score	Modelled data, NVIM Tool (DELWP 2017a)
	General Biodiversity Equivalence Score	= Habitat Hectares x Strategic Biodiversity Score
Moderate or High Risk-based pathway	Condition Score	Habitat hectare assessment
	Habitat Hectares	= Condition Score x Extent (ha)
	Strategic Biodiversity Score and Habitat Importance Score	Modelled data, determined by DEPI
	Specific Biodiversity Equivalence Score (A)	= Habitat Hectares x Habitat Importance Score
	Sum of Specific Biodiversity Equivalence Scores of remaining habitat (B)	Data gathered during the field assessment is provided to DEPI for analysis and a resulting assessment offset report is provided by the Department.
	Specific Offset Threshold (C)	
	General/Specific Threshold Test: If $A \div B > C$ a Specific offset is required If $A \div B < C$ a General offset required	

Table A1.5.2. Summary of offset requirements (DEPI 2013)

Risk-based Pathway	Offset Type	Offset Amount (Risk adjusted biodiversity equivalence score)	Habitat for Species	Offset Attributes	
				Vicinity	Strategic Biodiversity Score
Low Risk	General offset	1.5 times the general biodiversity equivalence score of the native vegetation to be removed.	No restrictions	In the same Catchment Management Authority or Local Government Area boundary as the native vegetation to be removed.	At least 80 per cent of the strategic biodiversity score of the native vegetation to be removed.
Moderate or High Risk	General offset	1.5 times the general biodiversity equivalence score of the native vegetation to be removed.	No restrictions	In the same Catchment Management Authority or Local Government Area boundary as the native vegetation to be removed.	At least 80 per cent of the strategic biodiversity score of the native vegetation to be removed.
Moderate or High Risk	Specific offset	For each species impacted, 2 times the specific biodiversity equivalence score of the native vegetation to be removed.	Likely habitat for each rare or threatened species that a specific offset is required for, according to the specific-general offset test.	No restrictions	No restrictions

Appendix 1.6 – Flora and Fauna Guarantee Act 1988 Protected Species

Protected flora and fauna under the *Flora and Fauna Guarantee Act 1988* (FFG Act) are defined as those that have legal protection under the Act. Protected taxa includes plants and animals from three sources:

- plant or animal taxa (species, subspecies or varieties) listed as threatened under the FFG Act;
- plant taxa belonging to communities listed as threatened under the FFG Act; and,
- plant taxa which are not threatened but require protection for other reasons.

Note that representative plants of a given community are protected as well as the community itself (for example scattered Wallaby-grasses *Rytidosperma* spp. are protected in degraded areas previously supporting the listed Western [Basalt] Plains Grassland Community).

Table A1.6 provides a list of plant groups protected under the FFG Act. For threatened plant species likely to occur within the study area refer to Appendix and for listed communities (or representative species) likely to occur within the study area refer to Sections 3.4.1 and 3.4.2.

Table A1.6. Plant groups (Families, Genera and Kingdom Divisions) protected under the FFG Act (DELWP 2016).

Family/Genera	Common Name	Exclusions
Pteridophyta	Clubmosses, ferns and fern allies	Austral Bracken <i>Pteridium esculentum</i>
Asteraceae	Daisies	N/A
Ericaceae (formerly Epacridaceae)	Heaths	N/A
Orchidaceae	Orchids	N/A
<i>Acacia</i>	Wattles	<i>Acacia dealbata</i> , <i>Acacia decurrens</i> , <i>Acacia implexa</i> , <i>Acacia melanoxylon</i> and <i>Acacia paradoxa</i>
<i>Baeckea</i>	Baeckeas	N/A
<i>Boronia</i>	Boronias	N/A
<i>Calytrix</i>	Fringe-myrtles	N/A
<i>Correa</i> -	Correas	N/A
<i>Darwinia</i>	Darwinias	N/A
<i>Eremophila</i>	Emu-bushes	N/A
<i>Eriostemon</i>	Wax-flowers	N/A
<i>Gompholobium</i>	Wedge-peas	N/A
<i>Grevillea</i>	Grevilleas	N/A
<i>Prostanthera</i>	Mint-bushes	N/A
<i>Sphagnum</i>	Sphagnum mosses	N/A
<i>Stylidium</i>	Trigger-plants	N/A
<i>Thryptomene</i>	Thryptomenes	N/A
<i>Thysanotus</i>	Fringe-lilies	N/A
<i>Xanthorrhoea</i>	Grass-trees	N/A

APPENDIX 2 - FLORA

Appendix 2.1 – Flora Results

Legend:

CR/EN/VU Listed as Critically Endangered/Endangered/Vulnerable under the EPBC Act;

I Protected under the FFG Act (DELWP 2018);

L Listed under the FFG Act (DELWP 2017e);

e/v/r/k Listed as endangered/vulnerable/rare/poorly known in Victoria under the Advisory List of Rare or Threatened Plants in Victoria (DEPI 2014);

***** Listed as a noxious weed under the CaLP Act;

w Weed of National Significance;

Planted Victorian and non-Victorian species;

+ Planted indigenous species that also occur in remnant native vegetation in the study area;

****** Planted indigenous species in the study area; and,

- Not applicable

Table A2.1. Flora recorded within the study area.

Scientific Name	Common Name	Comments
Indigenous Species		
<i>Acacia acinacea</i>	Gold-dust Wattle	I
<i>Acacia implexa</i>	Lightwood	
<i>Acacia paradoxa</i>	Hedge Wattle	
<i>Acacia pycnantha</i>	Golden Wattle	I
<i>Acacia rostriformis</i>	Bacchus Marsh Wattle	L I v **
<i>Acaena echinata</i>	Sheep's Burr	
<i>Asperula conferta</i>	Common Woodruff	
<i>Atriplex semibaccata</i>	Berry Saltbush	
<i>Austrostipa bigeniculata</i>	Kneed Spear-grass	
<i>Austrostipa densiflora</i>	Dense Spear-grass	
<i>Austrostipa elegantissima</i>	Feather Spear-grass	
<i>Austrostipa gibbosa</i>	Spurred Spear-grass	
<i>Austrostipa scabra</i> subsp. <i>falcata</i>	Rough Spear-grass	

Scientific Name	Common Name	Comments
<i>Brachyscome dentata</i>	Lobe-seed Daisy	l
<i>Calocephalus citreus</i>	Lemon Beauty-heads	l
<i>Carpobrotus modestus</i>	Inland Pigface	
<i>Cassinia arcuata</i>	Drooping Cassinia	l
<i>Centrolepis aristata</i>	Pointed Centrolepis	
<i>Chloris truncata</i>	Windmill Grass	
<i>Chrysocephalum semipapposum</i>	Clustered Everlasting	l
<i>Clematis microphylla</i> s.l.	Small-leaved Clematis	
<i>Convolvulus angustissimus</i> subsp. <i>omnigracilis</i>	Slender Bindweed	k
<i>Crassula decumbens</i> var. <i>decumbens</i>	Spreading Crassula	
<i>Crassula sieberiana</i> s.l.	Sieber Crassula	
<i>Dichondra repens</i>	Kidney-weed	
<i>Dodonaea viscosa</i>	Sticky Hop-bush	
<i>Einadia hastata</i>	Saloop	
<i>Einadia nutans</i>	Nodding Saltbush	
<i>Eleocharis acuta</i>	Common Spike-sedge	
<i>Enchylaena tomentosa</i> var. <i>tomentosa</i>	Ruby Saltbush	
<i>Eucalyptus behriana</i>	Bull Mallee	
<i>Eucalyptus leucoxylon</i> subsp. <i>connata</i>	Melbourne Yellow-gum	v
<i>Eucalyptus melliodora</i>	Yellow Box	
<i>Eucalyptus microcarpa</i>	Grey Box	
<i>Eucalyptus obliqua</i>	Messmate Stringybark	
<i>Goodenia ovata</i>	Hop Goodenia	
<i>Juncus holoschoenus</i>	Joint-leaf Rush	
<i>Juncus pallidus</i>	Pale Rush	
<i>Linum marginale</i>	Native Flax	
<i>Lomandra filiformis</i>	Wattle Mat-rush	
<i>Maireana enchylaenoides</i>	Wingless Bluebush	
<i>Melaleuca lanceolata</i>	Moonah	
<i>Melicytus denta</i>	Tree Violet	
<i>Oxalis perennans</i>	Grassland Wood-sorrel	
<i>Pimelea curviflora</i>	Curved Rice-flower	
<i>Pimelea spinescens</i> subsp. <i>spinescens</i>	Spiny Rice-flower	EN L e
<i>Pycnosorus chrysanthes</i>	Golden Billy-buttons	l
<i>Rhagodia parabolica</i>	Fragrant Saltbush	r
<i>Rumex brownii</i>	Slender Dock	
<i>Rytidosperma caespitosum</i>	Common Wallaby-grass	

Scientific Name	Common Name	Comments
<i>Rytidosperma geniculatum</i>	Knead Wallaby-grass	
<i>Rytidosperma setaceum</i>	Bristly Wallaby-grass	
<i>Sclerolaena diacantha</i>	Grey Copperburr	
<i>Sclerolaena muricata</i> var. <i>muricata</i>	Black Roly-poly	k
<i>Senecio pinnatifolius</i>	Variable Groundsel	l
<i>Senecio quadridentatus</i>	Cotton Fireweed	
<i>Themeda triandra</i>	Kangaroo Grass	
<i>Typha</i> spp.	Bulrush	
<i>Vittadinia cuneata</i>	Fuzzy New Holland Daisy	l
<i>Wahlenbergia gracilis</i>	Sprawling Bluebell	
<i>Wahlenbergia luteola</i>	Bronze Bluebell	
<i>Walwhalleya proluta</i>	Rigid Panic	
Introduced Species		
<i>Acetosella vulgaris</i>	Sheep Sorrel	
<i>Aira caryophyllea</i> subsp. <i>caryophyllea</i>	Silvery Hair-grass	
<i>Arctotheca calendula</i>	Cape Weed	
<i>Asparagus asparagoides</i>	Bridal Creeper	W *
<i>Avena</i> spp.	Oat	
<i>Brassica</i> spp.	Turnip	
<i>Briza minor</i>	Lesser Quaking-grass	
<i>Bromus catharticus</i>	Prairie Grass	
<i>Bromus diandrus</i>	Great Brome	
<i>Bromus hordeaceus</i> subsp. <i>hordeaceus</i>	Soft Brome	
<i>Cirsium vulgare</i>	Spear Thistle	*
<i>Conyza</i> spp.	Fleabane	
<i>Coprosma repens</i>	Mirror Bush	
<i>Cupressus macrocarpa</i>	Monterey Cypress	
<i>Cynara cardunculus</i> subsp. <i>flavescens</i>	Artichoke Thistle	*
<i>Cynodon dactylon</i> var. <i>dactylon</i>	Couch	
<i>Cyperus eragrostis</i>	Drain Flat-sedge	
<i>Dactylis glomerata</i>	Cocksfoot	
<i>Dittrichia graveolens</i>	Stinkwort	*
<i>Ehrharta erecta</i> var. <i>erecta</i>	Panic Veldt-grass	
<i>Eucalyptus botryoides</i>	Southern Mahogany	#
<i>Eucalyptus cladocalyx</i>	Sugar Gum	#
<i>Galenia pubescens</i> var. <i>pubescens</i>	Galenia	
<i>Gazania linearis</i>	Gazania	

Scientific Name	Common Name	Comments
<i>Helminthotheca echioides</i>	Ox-tongue	
<i>Holcus lanatus</i>	Yorkshire Fog	
<i>Hordeum</i> spp.	Barley	
<i>Hypochoeris radicata</i>	Flatweed	
<i>Lepidium africanum</i>	Common Peppergrass	
<i>Lolium perenne</i>	Perennial Rye-grass	
<i>Lycium ferocissimum</i>	African Box-thorn	W *
<i>Marrubium vulgare</i>	Horehound	W *
<i>Nassella neesiana</i>	Chilean Needle-grass	W *
<i>Nassella trichotoma</i>	Serrated Tussock	W *
<i>Olea europaea</i>	Olive	
<i>Opuntia</i> spp.	Prickly pear	W *
<i>Oxalis pes-caprae</i>	Soursob	*
<i>Paspalum dilatatum</i>	Paspalum	
<i>Phalaris aquatica</i>	Toowoomba Canary-grass	
<i>Physalis hederifolia</i>	Sticky Ground-cherry	
<i>Plantago coronopus</i>	Buck's-horn Plantain	
<i>Plantago lanceolata</i>	Ribwort	
<i>Prunus</i> spp.	Prunus	
<i>Romulea rosea</i>	Onion Grass	
<i>Rosa rubiginosa</i>	Sweet Briar	*
<i>Rubus fruticosus</i> spp. agg.	Blackberry	W *
<i>Rumex crispus</i>	Curled Dock	
<i>Salvia verbenaca</i>	Wild Sage	
<i>Schinus molle</i>	Pepper Tree	
<i>Solanum nigrum</i> s.l.	Black Nightshade	
<i>Sonchus asper</i> s.l.	Rough Sow-thistle	
<i>Sonchus oleraceus</i>	Common Sow-thistle	
<i>Tribulus terrestris</i>	Caltrop	
<i>Trifolium angustifolium</i> var. <i>angustifolium</i>	Narrow-leaf Clover	
<i>Trifolium</i> spp.	Clover	
<i>Vulpia bromoides</i>	Squirrel-tail Fescue	
<i>Vulpia myuros</i>	Rat's-tail Fescue	

Appendix 2.2 – Significant Flora Species

Table A2.2 Significant flora recorded within 10 kilometres of the study area

Likelihood: Habitat characteristics of significant flora species previously recorded within 10 kilometres of the study area, or that may potentially occur within the study area were assessed to determine their likelihood of occurrence. The likelihood of occurrence rankings are defined below.

1 - Known occurrence

- Recorded within the study area recently (i.e. within ten years)

2 - High Likelihood

- Previous records of the species in the local vicinity; and/or,
- The study area contains areas of high quality habitat.

3 - Moderate Likelihood

- Limited previous records of the species in the local vicinity; and/or,
- The study area contains poor or limited habitat.

4 - Low Likelihood

- Poor or limited habitat for the species however other evidence (such as a lack of records or environmental factors) indicates there is a very low likelihood of presence.

5 – Unlikely

- No suitable habitat and/or outside the species range.

Scientific name	Common name	Total # of documented records	Last documented record	EPBC	FFG	DEPI	Likely occurrence in study area
NATIONAL SIGNIFICANCE							
<i>Dianella amoena</i> #	Matted Flax-lily	-	-	EN	L	e	4
<i>Diuris basaltica</i>	Small Golden Moths	5	2012	EN	L	e	4
<i>Diuris fragrantissima</i>	Sunshine Diuris	1	1770	EN	L	e	4
<i>Eucalyptus aggregata</i> #	Black Gum	-	-	VU	L	e	4
<i>Glycine latrobeana</i> #	Clover Glycine	-	-	VU	L	v	3
<i>Lachnagrostis adamsonii</i> #	Adamson's Blown-grass	-	-	EN	L	v	4
<i>Leucochrysum albicans</i> var. <i>tricolor</i> #	Hoary Sunray	-	-	EN	-	e	4
<i>Pimelea spinescens</i> subsp. <i>spinescens</i>	Spiny Rice-flower	5	2003	CR	L	e	1
<i>Prasophyllum frenchii</i> #	Maroon Leek-orchid	-	-	EN	L	e	4

Scientific name	Common name	Total # of documented records	Last documented record	EPBC	FFG	DEPI	Likely occurrence in study area
<i>Senecio macrocarpus</i> #	Large-headed Fireweed	-	-	VU	L	e	3
<i>Thelymitra matthewsii</i> #	Spiral Sun-orchid	-	-	VU	L	v	4
<i>Xerochrysum palustre</i> #	Swamp Everlasting	-	-	VU	L	v	4
STATE SIGNIFICANCE							
<i>Acacia aspera</i> subsp. <i>parviceps</i>	Rough Wattle	2	1995	-	-	r	4
<i>Acacia rostriformis</i>	Bacchus Marsh Wattle	262	2016	-	L	v	1
<i>Allocasuarina luehmannii</i>	Buloke	15	2010	-	L	e	2
<i>Alternanthera</i> sp. 1 (Plains)	Plains Joyweed	2	2009	-	-	k	4
<i>Amyema linophylla</i> subsp. <i>orientalis</i>	Buloke Mistletoe	2	2010	-	-	v	4
<i>Austrostipa breviglumis</i>	Cane Spear-grass	18	2014	-	-	r	3
<i>Austrostipa exilis</i>	Heath Spear-grass	14	2008	-	-	r	3
<i>Boronia anemonifolia</i> subsp. <i>aurifodina</i>	Goldfield Boronia	1	1917	-	-	r	4
<i>Bossiaea cordigera</i>	Wiry Bossiaea	1	1980	-	-	r	4
<i>Calotis anthemoides</i>	Cut-leaf Burr-daisy	1	1984	-	L	-	3
<i>Calotis lappulacea</i>	Yellow Burr-daisy	2	1910	-	-	r	3
<i>Convolvulus angustissimus</i> subsp. <i>omnigracilis</i>	Slender Bindweed	5	2012	-	-	k	1
<i>Cullen parvum</i>	Small Scurf-pea	5	2012	-	L	e	2
<i>Cullen tenax</i>	Tough Scurf-pea	1	1853	-	L	e	3
<i>Desmodium varians</i>	Slender Tick-trefoil	3	2010	-	-	k	3
<i>Dianella</i> sp. aff. <i>longifolia</i> (Benambra)	Arching Flax-lily	7	2016	-	-	v	2
<i>Diuris gregaria</i>	Clumping Golden Moths	1	2012	-	L	e	3
<i>Eucalyptus</i> aff. <i>ignorabilis</i> (Lerderderg)	Lerderderg Scentbark	5	2011	-	-	e	3

Scientific name	Common name	Total # of documented records	Last documented record	EPBC	FFG	DEPI	Likely occurrence in study area
<i>Eucalyptus baueriana</i> subsp. <i>thalassina</i>	Werribee Blue-box	313	2011	-	-	e	2
<i>Eucalyptus leucoxydon</i> subsp. <i>connata</i>	Melbourne Yellow-gum	68	2016	-	-	v	2
<i>Euphrasia collina</i> subsp. <i>trichocalycina</i>	Purple Eyebright	1	1963	-	-	r	3
<i>Gahnia microstachya</i>	Slender Saw-sedge	6	2011	-	-	r	3
<i>Goodia medicaginea</i>	Western Golden-tip	2	1993	-	-	r	3
<i>Grevillea rosmarinifolia</i>	Rosemary Grevillea	2	1959	-	-	P	4
<i>Grevillea steiglitziana</i>	Brisbane Range Grevillea	1	1966	-	-	r	4
<i>Lepidium pseudohyssopifolium</i>	Native Peppergrass	2	2008	-	-	k	3
<i>Leucopogon microphyllus</i> var. <i>pilibundus</i>	Hairy Beard-heath	10	2011	-	-	r	3
<i>Maireana aphylla</i>	Leafless Bluebush	6	2006	-	-	k	3
<i>Marsilea mutica</i>	Smooth Nardoo	1	2000	-	-	k	3
<i>Myoporum montanum</i>	Waterbush	1	1853	-	-	r	3
<i>Nicotiana suaveolens</i>	Austral Tobacco	53	2013	-	-	r	2
<i>Olearia minor</i>	Satin Daisy-bush	1	1929	-	-	r	3
<i>Philothea angustifolia</i> subsp. <i>montana</i>	Narrow-leaf Wax-flower	1	1987	-	-	v	3
<i>Pimelea curviflora</i> var. aff. <i>subglabrata</i>	Curved Rice-flower	3	2001	-	-	k	3
<i>Pimelea hewardiana</i>	Forked Rice-flower	14	2011	-	-	r	2
<i>Poranthera corymbosa</i>	Clustered Poranthera	1	1982	-	-	r	3
<i>Prostanthera decussata</i>	Dense Mint-bush	1	1980	-	-	r	3
<i>Prostanthera nivea</i> var. <i>nivea</i>	Snowy Mint-bush	5	2011	-	-	r	3
<i>Prostanthera saxicola</i> var. <i>bracteolata</i>	Slender Mint-bush	1	2011	-	-	r	3
<i>Pseudanthus orbicularis</i>	Tangled Pseudanthus	3	1991	-	-	r	3

Scientific name	Common name	Total # of documented records	Last documented record	EPBC	FFG	DEPI	Likely occurrence in study area
<i>Pterostylis bicolor</i>	Black-tip Greenhood	1	1996	-	-	k	3
<i>Pterostylis truncata</i>	Brittle Greenhood	83	2013	-	L	e	3
<i>Ptilotus erubescens</i>	Hairy Tails	1	1984	-	L	v	3
<i>Pultenaea reflexifolia</i>	Wombat Bush-pea	1	1959	-	-	r	3
<i>Pultenaea weindorferi</i>	Swamp Bush-pea	1	1980	-	-	r	3
<i>Rhagodia parabolica</i>	Fragrant Saltbush	646	2016	-	-	r	1
<i>Sclerolaena muricata</i> var. <i>muricata</i>	Black Roly-poly	7	1998	-	-	k	1
<i>Senecio cunninghamii</i> var. <i>cunninghamii</i>	Branching Groundsel	5	2008	-	-	r	3
<i>Tripogon loliformis</i>	Rye Beetle-grass	2	2008	-	-	r	3
<i>Westringia glabra</i>	Violet Westringia	4	1980	-	-	r	3

Notes: EPBC = *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act), FFG = *Flora and Fauna Guarantee Act 1988* (FFG Act), DEPI= Advisory List of Rare or Threatened Plants in Victoria (DEPI 2014), L = Listed, # = Records identified from EPBC Act Protected Matters Search Tool, Data source: Victorian Biodiversity Atlas (DELWP 2017d); Protected Matters Search Tool (DoEE 2018). Order: Alphabetical. .

Appendix 2.3 – Habitat Hectare Results

Table A2.3. Habitat Hectares results for remnant vegetation recorded within the study area.

Vegetation Zone	GW ₁	GW ₂	GW ₃	GW ₄	PG ₁	PG ₂	PG ₃	PG ₄	PG ₅	PG ₆	PG ₇
Bioregion	CVU	CVU	CVU	CVU	VVP						
EVC / Tree	GW	GW	GW	GW	PG(LR)						
EVC Number	175	175	175	175	132_63	132_63	132_63	132_63	132_63	132_63	132_63
EVC Conservation Status	En	En	En	En	En	En	En	En	En	En	En
Patch Condition	Large Old Trees /10	7	7	0	5	0	0	0	0	0	0
	Canopy Cover /5	5	3	4	5	0	0	0	0	0	0
	Under storey /25	5	5	5	5	10	5	5	10	10	5
	Lack of Weeds /15	4	2	6	2	4	4	0	4	4	2
	Recruitment /10	3	1	1	3	3	1	0	3	0	6
	Organic Matter /5	5	5	0	3	4	2	2	5	4	5
	Logs /5	2	0	0	0	0	0	0	0	0	0
	Treeless EVC Multiplier	1.00	1.00	1.00	1.00	1.36	1.36	1.36	1.36	1.36	1.36
Subtotal =	31.00	23.00	16.00	23.00	28.56	16.32	9.52	29.92	24.48	24.48	36.72
Landscape Value /25	4	4	4	4	4	4	4	4	4	4	4
Habitat Points /100	35	27	20	27	33	20	14	34	28	28	41
Habitat Score	0.35	0.27	0.20	0.27	0.33	0.20	0.14	0.34	0.28	0.28	0.41
Total Area (ha)	1.586	0.109	0.043	3.748	0.341	3.926	6.783	3.727	2.174	1.258	0.663
Total habitat hectares	0.555	0.029	0.009	1.012	0.113	0.785	0.950	1.267	0.609	0.352	0.272
Conservation Significance	High	High	High	High	High	High	High	High	High	High	V. High
Large Old Trees in Patches	11	2	0	21	0	0	0	0	0	0	0

Vegetation Zone	PG8	PG9	PG10	RCW1	RCW2	RCW3	RCW4	RCW5	PGWe1	
Bioregion	VVP	VVP	VVP	CVU	CVU	CVU	CVU	CVU	CVU	
EVC / Tree	PG(LR)	PG(LR)	PG(LR)	RCW	RCW	RCW	RCW	RCW	PGWe	
EVC Number	132_63	132_63	132_63	64	64	64	64	64	125	
EVC Conservation Status	En	En	En	Vu	Vu	Vu	Vu	Vu	En	
Patch Condition	Large Old Trees /10	0	0	0	10	9	10	9	9	0
	Canopy Cover /5	0	0	0	4	4	5	5	5	0
	Under storey /25	15	10	10	5	10	15	15	15	10
	Lack of Weeds /15	9	6	2	2	4	11	9	0	4
	Recruitment /10	10	10	10	0	1	6	6	6	3
	Organic Matter /5	5	5	5	5	5	3	3	3	3
	Logs /5	0	0	0	0	0	4	4	4	0
	Treeless EVC Multiplier	1.36	1.36	1.36	1.00	1.00	1.00	1.00	1.00	1.36
Subtotal =	53.04	42.16	36.72	26.00	33.00	54.00	51.00	42.00	27.20	
Landscape Value /25	16	16	16	4	4	17	17	17	4	
Habitat Points /100	69	58	53	30	37	71	68	59	31	
Habitat Score	0.69	0.58	0.53	0.30	0.37	0.71	0.68	0.59	0.31	
Total Area (ha)	13.009	1.446	0.640	0.226	0.096	21.951	4.700	1.148	0.069	
Total habitat hectares	8.976	0.839	0.339	0.068	0.036	15.585	3.196	0.677	0.021	
Conservation Significance	V. High	V. High	V. High	High	High	V. High	V. High	V. High	High	
Large Old Trees in Patches	0	0	0	4	2	250	60	10	0	

Appendix 2.4 – Scattered Trees

Table A2.4. Remnant scattered trees recorded within the study area.

Tree ID	Common Name	Species Name	DBH	Size Class	Comments
1	Grey Box	<i>Eucalyptus microcarpa</i>	73	LOT	
2	Grey Box	<i>Eucalyptus microcarpa</i>	14	ST	
3	Grey Box	<i>Eucalyptus microcarpa</i>	34	ST	
4	Grey Box	<i>Eucalyptus microcarpa</i>	77	LOT	
5	Grey Box	<i>Eucalyptus microcarpa</i>	64	ST	Hollow
6	Grey Box	<i>Eucalyptus microcarpa</i>	62	ST	
7	Grey Box	<i>Eucalyptus microcarpa</i>	79	LOT	
8	Grey Box	<i>Eucalyptus microcarpa</i>	56	ST	Hollow
9	Grey Box	<i>Eucalyptus microcarpa</i>	76	LOT	
10	Grey Box	<i>Eucalyptus microcarpa</i>	102	LOT	Hollow
11	Grey Box	<i>Eucalyptus microcarpa</i>	97	LOT	
12	Grey Box	<i>Eucalyptus microcarpa</i>	87	LOT	
13	Grey Box	<i>Eucalyptus microcarpa</i>	79	LOT	
14	Grey Box	<i>Eucalyptus microcarpa</i>	78	LOT	
15	Grey Box	<i>Eucalyptus microcarpa</i>	110	LOT	
16	Grey Box	<i>Eucalyptus microcarpa</i>	105	LOT	Hollow
17	Grey Box	<i>Eucalyptus microcarpa</i>	41	ST	
18	Grey Box	<i>Eucalyptus microcarpa</i>	105	LOT	Hollow
19	Stag	Stag	74	LOT	Hollow
20	Messmate	<i>Eucalyptus obliqua</i>	110	LOT	
21	Grey Box	<i>Eucalyptus microcarpa</i>	88	LOT	
22	Grey Box	<i>Eucalyptus microcarpa</i>	112	LOT	Hollow
23	Grey Box	<i>Eucalyptus microcarpa</i>	9	ST	
24	Grey Box	<i>Eucalyptus microcarpa</i>	91	LOT	

Tree ID	Common Name	Species Name	DBH	Size Class	Comments
25	Grey Box	<i>Eucalyptus microcarpa</i>	88	LOT	
26	Grey Box	<i>Eucalyptus microcarpa</i>	14	ST	
27	Grey Box	<i>Eucalyptus microcarpa</i>	71	LOT	
28	Grey Box	<i>Eucalyptus microcarpa</i>	54	ST	
29	Grey Box	<i>Eucalyptus microcarpa</i>	61	ST	
30	Grey Box	<i>Eucalyptus microcarpa</i>	50	ST	
31	Grey Box	<i>Eucalyptus microcarpa</i>	65	ST	
32	Grey Box	<i>Eucalyptus microcarpa</i>	75	LOT	
33	Grey Box	<i>Eucalyptus microcarpa</i>	117	LOT	Hollow
34	River Red-gum	<i>Eucalyptus camaldulensis</i>	68	ST	
35	River Red-gum	<i>Eucalyptus camaldulensis</i>	59	ST	
36	River Red-gum	<i>Eucalyptus camaldulensis</i>	75	LOT	
37	Grey Box	<i>Eucalyptus microcarpa</i>	60	ST	
38	Grey Box	<i>Eucalyptus microcarpa</i>	92	LOT	Hollow
39	Grey Box	<i>Eucalyptus microcarpa</i>	106	LOT	Hollow
40	Grey Box	<i>Eucalyptus microcarpa</i>	95	LOT	Hollow
41	Yellow Box	<i>Eucalyptus melliodora</i>	21	ST	
42	River Red-gum	<i>Eucalyptus camaldulensis</i>	55	ST	
43	River Red-gum	<i>Eucalyptus camaldulensis</i>	61	LOT	
44	Stag	Stag	55	ST	
45	Yellow Box	<i>Eucalyptus melliodora</i>	17	ST	
46	Grey Box	<i>Eucalyptus microcarpa</i>	27	ST	
47	Grey Box	<i>Eucalyptus microcarpa</i>	25	ST	
48	Yellow Box	<i>Eucalyptus melliodora</i>	81	LOT	Hollow
49	River Red-gum	<i>Eucalyptus camaldulensis</i>	97	LOT	Hollow
50	Grey Box	<i>Eucalyptus microcarpa</i>	49	ST	
51	Grey Box	<i>Eucalyptus microcarpa</i>	71	LOT	

Tree ID	Common Name	Species Name	DBH	Size Class	Comments
52	Yellow Box	<i>Eucalyptus melliodora</i>	75	LOT	
53	Grey Box	<i>Eucalyptus microcarpa</i>	73	LOT	
54	Grey Box	<i>Eucalyptus microcarpa</i>	10	ST	
55	Grey Box	<i>Eucalyptus microcarpa</i>	78	LOT	Hollow
56	Yellow Box	<i>Eucalyptus melliodora</i>	55	ST	
57	Yellow Box	<i>Eucalyptus melliodora</i>	45	ST	
58	Grey Box	<i>Eucalyptus microcarpa</i>	37	ST	
59	Grey Box	<i>Eucalyptus microcarpa</i>	51	ST	
60	Stag	Stag	40	ST	

DRAFT

APPENDIX 3 - FAUNA

Appendix 3.1 – Significant Fauna Species

Table A3.1 Significant fauna within 10 kilometres of the study area.

Habitat characteristics of significant fauna species previously recorded within 10 kilometres of the study area, or that may potentially occur within the study area were assessed to determine their likelihood of occurrence. The likelihood of occurrence rankings for each of the threatened species are:

1	High Likelihood	<ul style="list-style-type: none"> • Known resident in the study area based on site observations, database records, or expert advice; and/or, • Recent records (i.e. within five years) of the species in the local area (VBA 2011); and/or, • The study area contains the species' preferred habitat.
2	Moderate Likelihood	<ul style="list-style-type: none"> • The species is likely to visit the study area regularly (i.e. at least seasonally); and/or, • Previous records of the species in the local area (DSE 2011b); and/or, • The study area contains some characteristics of the species' preferred habitat.
3	Low Likelihood	<ul style="list-style-type: none"> • The species is likely to visit the study area occasionally or opportunistically whilst en route to more suitable sites; and/or, • There are only limited or historical records of the species in the local area (i.e. more than 20 years old); and/or, • The study area contains few or no characteristics of the species' preferred habitat.
4	Unlikely	<ul style="list-style-type: none"> • No previous records of the species in the local area; and/or, • The species may fly over the study area when moving between areas of more suitable habitat; and/or, • Out of the species' range; and/or, • No suitable habitat present.

EX	Extinct
RX	Regionally extinct
CR	Critically endangered
EN	Endangered
VU	Vulnerable
RA	Rare
NT	Near threatened
CD	Conservation dependent
LC	least concern

DD	Data deficient (insufficiently or poorly known)
L	Listed as threatened under FFG Act
I	Invalid or ineligible for listing under the FFG Act
#	Listed on the Protected Matters Search Tool
*	Additional information from the Victorian Fauna Database

Common Name	Scientific Name	Last Documented Record (VBA)	# Records (VBA)	EPBC Act	FFG ACT	DSE (2013)	National Action Plan	Likelihood
NATIONAL SIGNIFICANCE								
Spot-tailed Quoll #	<i>Dasyurus maculatus maculatus</i>	-	-	EN	L	EN	VU	4
Eastern Barred Bandicoot	<i>Perameles gunnii</i>	1883	15	EN	L	WX	CR	4
Greater Glider #	<i>Petauroides volans</i>	-	-	VU	-	VU	VU	4
Grey-headed Flying-fox	<i>Pteropus poliocephalus</i>	1968	2	VU	L	VU	VU	3
Smoky Mouse #	<i>Pseudomys fumeus</i>	-	-	EN	L	EN	RA	4
Australasian Bittern	<i>Botaurus poiciloptilus</i>	1970	1	EN	L	EN	VU	4
Plains-wanderer	<i>Pedionomus torquatus</i>	1880	1	CR	L	CR	EN	3
Australian Painted Snipe	<i>Rostratula australis</i>	1989	1	VU	L	CR	VU	4
Eastern Curlew #	<i>Numenius madagascariensis</i>	-	-	CR	-	VU	-	4
Curlew Sandpiper #	<i>Calidris ferruginea</i>	-	-	CR	-	EN	-	4
Superb Parrot	<i>Polytelis swainsonii</i>	1881	1	VU	L	EN	VU	4
Swift Parrot	<i>Lathamus discolor</i>	2008	12	CR	L	EN	EN	2
Regent Honeyeater #	<i>Anthochaera phrygia</i>	-	-	CR	L	CR	EN	4
Painted Honeyeater #	<i>Grantiella picta</i>	-	-	VU	L	VU	NT	4
Pink-tailed Worm-Lizard #	<i>Aprasia parapulchella</i>	-	-	VU	L	EN	-	4
Striped Legless Lizard #	<i>Delma impar</i>	-	-	VU	L	EN	VU	2
Grassland Earless Dragon #	<i>Tympanocryptis pinguicolla</i>	-	-	EN	L	CR	VU	4
Growling Grass Frog	<i>Litoria raniformis</i>	2007	22	VU	L	EN	VU	4
Dwarf Galaxias #	<i>Galaxiella pusilla</i>	-	-	VU	L	EN	VU	4
Australian Grayling #	<i>Prototroctes maraena</i>	-	-	VU	L	VU	VU	4
Golden Sun Moth	<i>Synemon plana</i>	2012	333	CR	L	CR	-	1

Common Name	Scientific Name	Last Documented Record (VBA)	# Records (VBA)	EPBC Act	FFG ACT	DSE (2013)	National Action Plan	Likelihood
STATE SIGNIFICANCE								
Brush-tailed Phascogale	<i>Phascogale tapoatafa</i>	1989	9	-	L	VU	NT	3
Common Dunnart	<i>Sminthopsis murina murina</i>	1990	3	-	-	VU	-	4
Common Bent-wing Bat	<i>Miniopterus schreibersii</i> GROUP	1999	5	-	L	-	CD	3
Musk Duck	<i>Biziura lobata</i>	2003	19	-	-	VU	-	4
Freckled Duck	<i>Stictonetta naevosa</i>	2006	6	-	L	EN	-	4
Australasian Shoveler	<i>Anas rhynchotis</i>	2006	11	-	-	VU	-	4
Hardhead	<i>Aythya australis</i>	2006	16	-	-	VU	-	4
Blue-billed Duck	<i>Oxyura australis</i>	2006	4	-	L	EN	-	4
Diamond Dove	<i>Geopelia cuneata</i>	1905	2	-	L	NT	-	4
White-throated Needletail	<i>Hirundapus caudacutus</i>	1994	9	-	-	VU	-	3
Eastern Great Egret	<i>Ardea modesta</i>	2001	8	-	L	VU	-	3
Little Egret	<i>Egretta garzetta nigripes</i>	1990	1	-	L	EN	-	3
White-bellied Sea-Eagle	<i>Haliaeetus leucogaster</i>	2006	5	-	L	VU	-	4
Black Falcon	<i>Falco subniger</i>	1988	8	-	-	VU	-	3
Lewin's Rail	<i>Lewinia pectoralis pectoralis</i>	1880	1	-	L	VU	NT	4
Australian Bustard	<i>Ardeotis australis</i>	1911	1	-	L	CR	NT	4
Bush Stone-curlew	<i>Burhinus grallarius</i>	1880	1	-	L	EN	NT	4
Gull-billed Tern	<i>Gelochelidon nilotica macrotarsa</i>	1986	1	-	L	EN	-	4
Caspian Tern	<i>Hydroprogne caspia</i>	2000	1	-	L	NT	-	4
Powerful Owl	<i>Ninox strenua</i>	2011	10	-	L	VU	-	3
Barking Owl	<i>Ninox connivens connivens</i>	2002	25	-	L	EN	NT	3

Common Name	Scientific Name	Last Documented Record (VBA)	# Records (VBA)	EPBC Act	FFG ACT	DSE (2013)	National Action Plan	Likelihood
Brown Treecreeper (south-eastern ssp.)	<i>Climacteris picumnus victoriae</i>	2010	75	-	-	NT	NT	2
Chestnut-rumped Heathwren	<i>Calamanthus pyrrhopygius</i>	2004	10	-	L	VU	-	4
Speckled Warbler	<i>Chthonicola sagittatus</i>	2014	82	-	L	VU	NT	2
Grey-crowned Babbler	<i>Pomatostomus temporalis temporalis</i>	1880	1	-	L	EN	NT	4
Crested Bellbird	<i>Oreoica gutturalis gutturalis</i>	2003	18	-	L	NT	NT	3
Hooded Robin	<i>Melanodryas cucullata cucullata</i>	1999	12	-	L	NT	NT	3
Diamond Firetail	<i>Stagonopleura guttata</i>	2011	69	-	L	NT	NT	2
Bearded Dragon	<i>Pogona barbata</i>	1986	2	-	-	VU	-	4
Lace Goanna	<i>Varanus varius</i>	1987	2	-	-	EN	-	3
Brown Toadlet	<i>Pseudophryne bibronii</i>	1990	5	-	L	EN	DD	4
Bullant	<i>Myrmecia</i> sp. 17	2009	4	-	L	VU	-	2
REGIONAL SIGNIFICANCE								
Fat-tailed Dunnart	<i>Sminthopsis crassicaudata</i>	1988	1	-	-	NT	-	2
Eastern Pygmy-possum	<i>Cercartetus nanus</i>	1992	2	-	-	NT	-	4
Pied Cormorant	<i>Phalacrocorax varius</i>	2003	4	-	-	NT	-	3
Nankeen Night Heron	<i>Nycticorax caledonicus hillii</i>	2000	11	-	-	NT	-	3
Glossy Ibis	<i>Plegadis falcinellus</i>	1986	2	-	-	NT	-	3
Royal Spoonbill	<i>Platalea regia</i>	1991	6	-	-	NT	-	3
Spotted Harrier	<i>Circus assimilis</i>	2008	7	-	-	NT	-	2
Latham's Snipe	<i>Gallinago hardwickii</i>	1990	5	-	-	NT	-	3
Little Button-quail	<i>Turnix velox</i>	2011	2	-	-	NT	-	3
Black-eared Cuckoo	<i>Chrysococcyx osculans</i>	2000	13	-	-	NT	-	4

Common Name	Scientific Name	Last Documented Record (VBA)	# Records (VBA)	EPBC Act	FFG ACT	DSE (2013)	National Action Plan	Likelihood
Azure Kingfisher	<i>Alcedo azurea</i>	1988	2	-	-	NT	-	4
Red-backed Kingfisher	<i>Todiramphus pyrropygia pyrropygia</i>	1986	1	-	-	NT	-	4
Spotted Quail-thrush	<i>Cinlosoma punctatum</i>	2010	4	-	-	NT	-	3

EPBC *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act)

FFG *Flora and Fauna Guarantee Act 1988* (FFG Act)

DSE Advisory List of Threatened Vertebrate Fauna in Victoria (DSE 2013); Advisory List of Threatened Invertebrate Fauna in Victoria (DSE 2009)

NAP National Action Plan (Cogger *et al.* 1993; Duncan *et al.* 1999; Garnet and Crowley 2000; Lee 1995; Maxwell *et al.* 1996; Sands and New 2002; Tyler 1997)

Data source: Victorian Biodiversity Atlas (DELWP 2017e); Protected Matters Search Tool (DoEE 2018).

Taxonomic order: Mammals (Strahan 1995 *in* Menkhorst and Knight 2004); Birds (Christidis and Boles, 2008); Reptiles and Amphibians (Cogger *et al.* 1983 *in* Cogger 1996); Fish (Nelson 1994).

APPENDIX 4 - NATIVE VEGETATION REPORTS

Appendix 4.1 – Native Vegetation Removal Report

DRAFT

This report provides information to support an application to remove, destroy or lop native vegetation in accordance with the *Guidelines for the removal, destruction or lopping of native vegetation*. The report **is not an assessment by DELWP** of the proposed native vegetation removal. Native vegetation information and offset requirements have been determined using spatial data provided by the applicant or their consultant.

Date of issue: 02/07/2018

Report ID: EHP_2018_172


Time of issue: 12:00 pm

Project ID	EHP10937_Bacchus_Marsh
------------	------------------------

Assessment pathway

Assessment pathway	Detailed Assessment Pathway
Extent including past and proposed	12.069 ha
Extent of past removal	0.000 ha
Extent of proposed removal	12.069 ha
No. Large trees proposed to be removed	0
Location category of proposed removal	Location 3 The native vegetation is in an area where the removal of less than 0.5 hectares could have a significant impact on habitat for one or more rare or threatened species. The native vegetation is also in an area mapped as an endangered Ecological Vegetation Class (as per the statewide EVC map).

1. Location map



Offset requirements if a permit is granted

Any approval granted will include a condition to obtain an offset that meets the following requirements:

General offset amount¹	0.001 general habitat units
Vicinity	Port Phillip and Westernport Catchment Management Authority (CMA) or Moorabool Shire Council
Minimum strategic biodiversity value score ²	0.800
Large trees*	0 large trees
Species offset amount³	9.744 species units of habitat for Small Golden Moths, <i>Diuris basaltica</i> 10.049 species units of habitat for Heath Spear-grass, <i>Austrostipa exilis</i> 10.108 species units of habitat for Melbourne Yellow-gum, <i>Eucalyptus leucoxylon subsp. connata</i> 9.675 species units of habitat for Basalt Podolepis, <i>Podolepis linearifolia</i> 10.079 species units of habitat for Bacchus Marsh Wattle, <i>Acacia rostriformis</i> 10.049 species units of habitat for Fragrant Saltbush, <i>Rhagodia parabolica</i>
Large trees*	0 trees
* The total number of large trees that the offset must protect	0 large trees to be protected in either the general, species or combination across all habitat units protected

NB: values within tables in this document may not add to the totals shown above due to rounding

Appendix 1 includes information about the native vegetation to be removed

Appendix 2 includes information about the rare or threatened species mapped at the site.

Appendix 3 includes maps showing native vegetation to be removed and extracts of relevant species habitat importance maps

¹ The general offset amount required is the sum of all general habitat units in Appendix 1.

² Minimum strategic biodiversity score is 80 per cent of the weighted average score across habitat zones where a general offset is required

³ The species offset amount(s) required is the sum of all species habitat units in Appendix 1.

Next steps

Any proposal to remove native vegetation must meet the application requirements of the Detailed Assessment Pathway and it will be assessed under the Detailed Assessment Pathway.

If you wish to remove the mapped native vegetation you are required to apply for a permit from your local council. Council will refer your application to DELWP for assessment, as required. **This report is not a referral assessment by DELWP.**

This *Native vegetation removal report* must be submitted with your application for a permit to remove, destroy or lop native vegetation.

Refer to the *Guidelines for the removal, destruction or lopping of native vegetation* (the Guidelines) for a full list of application requirements. This report provides information that meets the following application requirements:

- The assessment pathway and reason for the assessment pathway
- A description of the native vegetation to be removed (partly met)
- Maps showing the native vegetation and property (partly met)
- Information about the impacts on rare or threatened species.
- The offset requirements determined in accordance with section 5 of the Guidelines that apply if approval is granted to remove native vegetation.

Additional application requirements must be met including:

- Topographical and land information
- Recent dated photographs
- Details of past native vegetation removal
- An avoid and minimise statement
- A copy of any Property Vegetation Plan that applies
- A defensible space statement as applicable
- A statement about the Native Vegetation Precinct Plan as applicable
- A site assessment report including a habitat hectare assessment of any patches of native vegetation and details of trees
- An offset statement that explains that an offset has been identified and how it will be secured.

© The State of Victoria Department of Environment, Land, Water and Planning Melbourne 2018

This work is licensed under a Creative Commons Attribution 4.0 International licence. You are free to re-use the work under that licence, on the condition that you credit the State of Victoria as author. The licence does not apply to any images, photographs or branding, including the Victorian Coat of Arms, the Victorian Government logo and the Department of Environment, Land, Water and Planning logo. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/3.0/au/deed.en>

Authorised by the Victorian Government, 8 Nicholson Street, East Melbourne.

For more information contact the DELWP Customer Service Centre 136 186

www.delwp.vic.gov.au

Disclaimer

This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

Obtaining this publication does not guarantee that an application will meet the requirements of Clauses 52.16 or 52.17 of the Victoria Planning Provisions and Victorian planning schemes or that a permit to remove native vegetation will be granted.

Notwithstanding anything else contained in this publication, you must ensure that you comply with all relevant laws, legislation, awards or orders and that you obtain and comply with all permits, approvals and the like that affect, are applicable or are necessary to undertake any action to remove, lop or destroy or otherwise deal with any native vegetation or that apply to matters within the scope of Clauses 52.16 or 52.17 of the Victoria Planning Provisions and Victorian planning schemes.

Appendix 1: Description of native vegetation to be removed

The species-general offset test was applied to your proposal. This test determines if the proposed removal of native vegetation has a proportional impact on any rare or threatened species habitats above the species offset threshold. The threshold is set at 0.005 per cent of the mapped habitat value for a species. When the proportional impact is above the species offset threshold a species offset is required. This test is done for all species mapped at the site. Multiple species offsets will be required if the species offset threshold is exceeded for multiple species.

Where a zone requires species offset(s), the species habitat units for each species in that zone is calculated by the following equation in accordance with the Guidelines:

$$\text{Species habitat units} = \text{extent} \times \text{condition} \times \text{species landscape factor} \times 2, \text{ where the species landscape factor} = 0.5 + (\text{habitat importance score}/2)$$

The species offset amount(s) required is the sum of all species habitat units per zone

Where a zone does not require a species offset, the general habitat units in that zone is calculated by the following equation in accordance with the Guidelines:

$$\text{General habitat units} = \text{extent} \times \text{condition} \times \text{general landscape factor} \times 1.5, \text{ where the general landscape factor} = 0.5 + (\text{strategic biodiversity value score}/2)$$

The general offset amount required is the sum of all general habitat units per zone.

Native vegetation to be removed

Zone	Type	BioEVC	BioEVC conservation status	Large tree(s)	Partial removal	Information provided by or on behalf of the applicant in a GIS file			Information calculated by EnSym					
						Condition score	Polygon Extent	Extent without overlap	SBV score	HI score	Habitat units	Offset type		
1-G	Patch	vp_0132_61	Endangered	0	no	0.200	2.263	2.263	0.925	0.726	0.781	501473	Small Golden Moths	<i>Diuris basaltica</i>
										0.499	0.781	503984	Heath Spear-grass	<i>Austrostipa exilis</i>
										0.499	0.781	504484	Melbourne Yellow-gum	<i>Eucalyptus leucoxylo</i> subsp. <i>connata</i>
										0.499	0.781	504658	Basalt Podolepis	<i>Podolepis linearifolia</i>
										0.439	0.781	505136	Bacchus Marsh Wattle	<i>Acacia rostriformis</i>
										0.499	0.781	502929	Fragrant Saltbush	<i>Rhagodia parabolica</i>
2-G	Patch	vp_0132_61	Endangered	0	no	0.200	0.055	0.055	0.852	0.743	0.019	501473	Small Golden Moths	<i>Diuris basaltica</i>
										0.735	0.019	503984	Heath Spear-grass	<i>Austrostipa exilis</i>
										0.735	0.019	504484	Melbourne Yellow-gum	<i>Eucalyptus leucoxylo</i> subsp. <i>connata</i>

Information provided by or on behalf of the applicant in a GIS file							Information calculated by EnSym					
Zone	Type	BioEVC	BioEVC conservation status	Large tree(s)	Partial removal	Condition score	Polygon Extent	Extent without overlap	SBV score	HI score	Habitat units	Offset type
										0.735	0.019	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.735	0.019	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
3-G	Patch	wp_0132_61	Endangered	0	no	0.200	0.018	0.018	0.830	0.720	0.006	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.720	0.006	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxyton subsp. connata</i>
										0.720	0.006	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.720	0.006	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
4-I	Patch	wp_0132_61	Endangered	0	no	0.340	1.782	1.782	0.547	0.683	1.020	501473 Small Golden Moths <i>Diuris basaltica</i>
										0.254	1.019	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.254	1.019	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxyton subsp. connata</i>
										0.103	1.006	504658 Basalt Podolepis <i>Podolepis linearifolia</i>
										0.254	1.019	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.254	1.019	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
5-H	Patch	wp_0132_61	Endangered	0	no	0.140	0.659	0.659	0.807	0.728	0.159	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.728	0.159	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxyton subsp. connata</i>
										0.728	0.159	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.728	0.159	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
6-J	Patch	wp_0132_61	Endangered	0	no	0.280	0.043	0.043	0.770	0.810	0.022	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.810	0.022	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxyton subsp. connata</i>

Information provided by or on behalf of the applicant in a GIS file							Information calculated by EnSym					
Zone	Type	BioEVC	BioEVC conservation status	Large tree(s)	Partial removal	Condition score	Polygon Extent	Extent without overlap	SBV score	HI score	Habitat units	Offset type
										0.810	0.022	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.810	0.022	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
7-G	Patch	wp_0132_61	Endangered	0	no	0.200	0.560	0.560	0.900	0.718	0.192	501473 Small Golden Moths <i>Diuris basaltica</i>
										0.718	0.192	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.718	0.192	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxydon subsp. connata</i>
										0.398	0.193	504658 Basalt Podolepis <i>Podolepis linearifolia</i>
										0.667	0.192	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.718	0.192	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
8-G	Patch	wp_0132_61	Endangered	0	no	0.200	0.816	0.816	0.865	0.710	0.279	501473 Small Golden Moths <i>Diuris basaltica</i>
										0.641	0.268	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.641	0.268	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxydon subsp. connata</i>
										0.147	0.243	504658 Basalt Podolepis <i>Podolepis linearifolia</i>
										0.641	0.268	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.641	0.268	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
9-G	Patch	wp_0132_61	Endangered	0	no	0.200	0.014	0.014	0.890	0.780	0.005	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.780	0.005	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxydon subsp. connata</i>
										0.780	0.005	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.780	0.005	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
10-H	Patch	wp_0132_61	Endangered	0	no	0.140	0.364	0.364	0.437	0.710	0.087	503984 Heath Spear-grass <i>Austrostipa exilis</i>

Information provided by or on behalf of the applicant in a GIS file							Information calculated by EnSym					
Zone	Type	BioEVC	BioEVC conservation status	Large tree(s)	Partial removal	Condition score	Polygon Extent	Extent without overlap	SBV score	HI score	Habitat units	Offset type
										0.407	0.087	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxylo</i> subsp. <i>connata</i>
										0.407	0.087	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.407	0.087	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
11-H	Patch	wp_0132_61	Endangered	0	no	0.140	0.116	0.116	0.820	0.406	0.023	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.406	0.023	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxylo</i> subsp. <i>connata</i>
										0.406	0.023	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.406	0.023	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
12-H	Patch	wp_0132_61	Endangered	0	no	0.140	0.058	0.058	0.820	0.409	0.011	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.409	0.011	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxylo</i> subsp. <i>connata</i>
										0.409	0.011	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.409	0.011	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
13-U	Patch	wp_0132_61	Endangered	0	no	0.785	5.320	5.320	0.960	0.784	7.452	501473 Small Golden Moths <i>Diuris basaltica</i>
										0.785	7.456	503984 Heath Spear-grass <i>Austrostipa exilis</i>
										0.593	7.515	504484 Melbourne Yellow-gum <i>Eucalyptus leucoxylo</i> subsp. <i>connata</i>
										0.733	7.452	504658 Basalt Podolepis <i>Podolepis linearifolia</i>
										0.428	7.486	505136 Bacchus Marsh Wattle <i>Acacia rostriformis</i>
										0.785	7.456	502929 Fragrant Saltbush <i>Rhagodia parabolica</i>
17-E	Patch	wp_0132_61	Endangered	0	no	0.330	0.002	0.002	1.000		0.001	General

Appendix 2: Information about impacts to rare or threatened species' habitats on site

This table lists all rare or threatened species' habitats mapped at the site.

Species common name	Species scientific name	Species number	Conservation status	Group	Habitat impacted	% habitat value affected
Small Golden Moths	<i>Diuris basaltica</i>	501473	Endangered	Dispersed	Top ranking map	0.0426
Heath Spear-grass	<i>Austrostipa exilis</i>	503984	Rare	Dispersed	Top ranking map	0.0393
Bacchus Marsh Wattle	<i>Acacia rostriformis</i>	505136	Vulnerable	Dispersed	Top ranking map	0.0372
Basalt Podolepis	<i>Podolepis linearifolia</i>	504658	Endangered	Dispersed	Top ranking map	0.0303
Melbourne Yellow-gum	<i>Eucalyptus leucoxylon subsp. connata</i>	504484	Vulnerable	Dispersed	Top ranking map	0.0211
Bacchus Marsh Wattle	<i>Acacia rostriformis</i>	505136	Vulnerable	Dispersed	Habitat importance map	0.0137
Small Golden Moths	<i>Diuris basaltica</i>	501473	Endangered	Dispersed	Habitat importance map	0.0093
Heath Spear-grass	<i>Austrostipa exilis</i>	503984	Rare	Dispersed	Habitat importance map	0.0088
Fragrant Saltbush	<i>Rhagodia parabolica</i>	502929	Rare	Dispersed	Habitat importance map ; special site	0.0075
Melbourne Yellow-gum	<i>Eucalyptus leucoxylon subsp. connata</i>	504484	Vulnerable	Dispersed	Habitat importance map	0.0041
Basalt Podolepis	<i>Podolepis linearifolia</i>	504658	Endangered	Dispersed	Habitat importance map	0.0037
Button Wrinklewort	<i>Rutidosis leptorhynchoides</i>	502982	Endangered	Dispersed	Habitat importance map	0.0032
Grassland Earless Dragon	<i>Tympanocryptis pinguicolla</i>	12922	Critically endangered	Dispersed	Habitat importance map	0.0029
Large-headed Fireweed	<i>Senecio macrocarpus</i>	503116	Endangered	Dispersed	Habitat importance map	0.0024
Large-flower Crane's-bill	<i>Geranium sp. 1</i>	505342	Endangered	Dispersed	Habitat importance map	0.0022
Plump Swamp Wallaby-grass	<i>Amphibromus pithogastrus</i>	503624	Endangered	Dispersed	Habitat importance map	0.0020
Austral Tobacco	<i>Nicotiana suaveolens</i>	502275	Rare	Dispersed	Habitat importance map ; special site	0.0020
Velvet Daisy-bush	<i>Olearia pannosa subsp. cardiophylla</i>	502317	Vulnerable	Dispersed	Habitat importance map	0.0020

Brackish Plains Buttercup	<i>Ranunculus diminutus</i>	504314	Rare	Dispersed	Habitat importance map	0.0019
Cane Spear-grass	<i>Austrostipa breviglumis</i>	503268	Rare	Dispersed	Habitat importance map	0.0015
Small Scurf-pea	<i>Cullen parvum</i>	502773	Endangered	Dispersed	Habitat importance map	0.0015
Yellow Burr-daisy	<i>Calotis lappulacea</i>	500598	Rare	Dispersed	Habitat importance map	0.0013
Snowy Mint-bush	<i>Prostanthera nivea</i> var. <i>nivea</i>	502746	Rare	Dispersed	Habitat importance map	0.0013
Spiny Rice-flower	<i>Pimelea spinescens</i> subsp. <i>spinescens</i>	504823	Endangered	Dispersed	Habitat importance map	0.0012
Clumping Golden Moths	<i>Diuris gregaria</i>	504887	Endangered	Dispersed	Habitat importance map	0.0012
Matted Flax-lily	<i>Dianella amoena</i>	505084	Endangered	Dispersed	Habitat importance map	0.0011
Tough Scurf-pea	<i>Cullen tenax</i>	502776	Endangered	Dispersed	Habitat importance map	0.0011
Rye Beetle-grass	<i>Tripogon loliformis</i>	503455	Rare	Dispersed	Habitat importance map	0.0011
Pale-flower Crane's-bill	<i>Geranium</i> sp. 3	505344	Rare	Dispersed	Habitat importance map	0.0011
Dark Wire-grass	<i>Aristida calycina</i> var. <i>calycina</i>	503630	Rare	Dispersed	Habitat importance map	0.0009
Arching Flax-lily	<i>Dianella</i> sp. aff. <i>longifolia</i> (<i>Benambra</i>)	505560	Vulnerable	Dispersed	Habitat importance map	0.0009
Pale Swamp Everlasting	<i>Coronidium gunnianum</i>	504655	Vulnerable	Dispersed	Habitat importance map	0.0008
Golden Sun Moth	<i>Synemon plana</i>	15021	Critically endangered	Dispersed	Habitat importance map	0.0008
Rosemary Grevillea	<i>Grevillea rosmarinifolia</i> subsp. <i>rosmarinifolia</i>	504066	Rare	Dispersed	Habitat importance map	0.0008
Late-flower Flax-lily	<i>Dianella tarda</i>	505085	Vulnerable	Dispersed	Habitat importance map	0.0007
Fragrant Saltbush	<i>Rhagodia parabolica</i>	502929	Rare	Dispersed	Top ranking map ; special site	0.0005
Small Milkwort	<i>Comesperma polygaloides</i>	500798	Vulnerable	Dispersed	Habitat importance map	0.0005
Purple Diuris	<i>Diuris punctata</i>	501084	Vulnerable	Dispersed	Habitat importance map	0.0005
Button Wrinklewort	<i>Rutidosis leptorhynchoides</i>	502982	Endangered	Dispersed	Top ranking map	0.0005
Waterbush	<i>Myoporum montanum</i>	502240	Rare	Dispersed	Habitat importance map	0.0004
Hairy Tails	<i>Ptilotus erubescens</i>	502825	Vulnerable	Dispersed	Habitat importance map	0.0004

Brittle Greenhood	<i>Pterostylis truncata</i>	502821	Endangered	Dispersed	Habitat importance map	0.0004
Clover Glycine	<i>Glycine latrobeana</i>	501456	Vulnerable	Dispersed	Habitat importance map	0.0003
Golden Cowslips	<i>Diuris behrii</i>	501061	Vulnerable	Dispersed	Habitat importance map	0.0003
Large-headed Fireweed	<i>Senecio macrocarpus</i>	503116	Endangered	Dispersed	Top ranking map	0.0003
Branching Groundsel	<i>Senecio cunninghamii</i> var. <i>cunninghamii</i>	503104	Rare	Dispersed	Habitat importance map	0.0002
Buloke	<i>Allocasuarina luehmannii</i>	500678	Endangered	Dispersed	Habitat importance map ; special site	0.0002
Black Falcon	<i>Falco subniger</i>	10238	Vulnerable	Dispersed	Habitat importance map	0.0002
Shiny Leionema	<i>Leionema lamprophyllum</i> subsp. <i>obovatum</i>	505478	Rare	Dispersed	Habitat importance map	0.0002
Port Lincoln Snake	<i>Parasuta spectabilis</i>	12813	Vulnerable	Dispersed	Habitat importance map	0.0001
Austral Crane's-bill	<i>Geranium solanderi</i> var. <i>solanderi</i> s.s.	505337	Vulnerable	Dispersed	Habitat importance map	0.0001
Forked Rice-flower	<i>Pimelea hewardiana</i>	502522	Rare	Dispersed	Habitat importance map	0.0001
Dwarf Brooklime	<i>Gratiola pumilo</i>	503753	Rare	Dispersed	Habitat importance map	0.0000
Western Golden-tip	<i>Goodia medicaginea</i>	501518	Rare	Dispersed	Habitat importance map	0.0000
Hairy Beard-heath	<i>Leucopogon microphyllus</i> var. <i>pilibundus</i>	501988	Rare	Dispersed	Habitat importance map	0.0000
Trailing Hop-bush	<i>Dodonaea procumbens</i>	501090	Vulnerable	Dispersed	Habitat importance map	0.0000
Dense Mint-bush	<i>Prostanthera decussata</i>	502739	Rare	Dispersed	Habitat importance map	0.0000
Striped Legless Lizard	<i>Delma impar</i>	12159	Endangered	Dispersed	Habitat importance map	0.0000
Speckled Warbler	<i>Chthonicola sagittatus</i>	10504	Vulnerable	Dispersed	Habitat importance map	0.0000
Silky Kidney-weed	<i>Dichondra</i> sp. 1	505786	Rare	Dispersed	Habitat importance map	0.0000
Brown Toadlet	<i>Pseudophryne bibronii</i>	13117	Endangered	Dispersed	Habitat importance map	0.0000
Half-bearded Spear-grass	<i>Austrostipa hemipogon</i>	503985	Rare	Dispersed	Habitat importance map	0.0000
Common Dunnart	<i>Sminthopsis murina murina</i>	11061	Vulnerable	Dispersed	Habitat importance map	0.0000
Barking Owl	<i>Ninox connivens connivens</i>	10246	Endangered	Dispersed	Habitat importance map	0.0000

Chestnut-rumped Heathwren	<i>Calamanthus pyrrhopygius</i>	10498	Vulnerable	Dispersed	Habitat importance map	0.0000
Lace Monitor	<i>Varanus varius</i>	12283	Endangered	Dispersed	Habitat importance map	0.0000
Slender Mint-bush	<i>Prostanthera saxicola var. bracteolata</i>	502750	Rare	Dispersed	Habitat importance map	0.0000


Habitat group

- Highly localised habitat means there is 2000 hectares or less mapped habitat for the species
- Dispersed habitat means there is more than 2000 hectares of mapped habitat for the species

Habitat impacted

- Habitat importance maps are the maps defined in the Guidelines that include all the mapped habitat for a rare or threatened species
- Top ranking maps are the maps defined in the Guidelines that depict the important areas of a dispersed species habitat, developed from the highest habitat importance scores in dispersed species habitat maps and selected VBA records
- Selected VBA record is an area in Victoria that represents a large population, roosting or breeding site etc.



4. Map of the property in context





Yellow boundaries denote areas of proposed native vegetation removal.

4. Habitat importance maps



Small Golden Moths
Diuris basaltica
501473





Heath Spear-grass
Austrostipa exilis
503984




Melbourne Yellow-gum
Eucalyptus leucoxylon subsp. connata
504484




Basalt Podolepis
Podolepis linearifolia
504658



Bacchus Marsh Wattle
Acacia rostriformis
505136



Fragrant Saltbush
Rhagodia parabolica
502929



Appendix 4.2 – Native Vegetation Offset Report

DRAFT

Native vegetation offset report

Information included in this report is based on spatial data provided to DELWP. The proposal has not been assessed to confirm eligibility or gain

This report provides information about a potential native vegetation offset site in accordance with the *Guidelines for the removal, destruction or lopping of native vegetation*. The information in this report is based on spatial information and the gain score provided by the landholder (or their representative). Any changes to this input information will change the habitat units of gain reflected in this report and it must be reissued.

Date of issue: 02/08/2018
Time of issue: 1:57 pm

DELWP ref: EHP_2018_0203

Project ID	EHP10937_Bacchus_Marsh
------------	------------------------

Extent of proposed offset site

Total extent	42.869 ha
Patches	42.869 ha
Revegetation	0.000 ha
Scattered tree(s)	0.000 ha

Habitat units of gain for the proposed offset site

The offset site has the following total general and species habitat units. These units can be used to satisfy a **single permit condition** or if the offset site is established as a **first party offset site**.

Total habitat units and attributes used for a single permit (once off use)

Number of large tree(s)	11 large trees are protected at the offset site
General habitat units	12.973 general habitat units Port Phillip And Westernport CMA, Moorabool Shire Council 0.895 Strategic biodiversity value
Species habitat units	12.114 species habitat units for Bacchus Marsh Wattle, <i>Acacia rostriformis</i> 5.558 species habitat units for Werribee Blue-box, <i>Eucalyptus baueriana subsp. thalassina</i> 6.309 species habitat units for Swift Parrot, <i>Lathamus discolor</i> 9.867 species habitat units for Speckled Warbler, <i>Chthonicola sagittatus</i> 12.213 species habitat units for Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i> 7.627 species habitat units for Brown Toadlet, <i>Pseudophryne bibronii</i> 12.956 species habitat units for Golden Sun Moth, <i>Synemon plana</i>

Native vegetation offset report

12.118 species habitat units for Yellow Burr-daisy, <i>Calotis lappulacea</i>
11.970 species habitat units for Trailing Hop-bush, <i>Dodonaea procumbens</i>
4.757 species habitat units for Small Golden Moths, <i>Diuris basaltica</i>
11.939 species habitat units for Western Golden-tip, <i>Goodia medicaginea</i>
11.944 species habitat units for Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
12.109 species habitat units for Austral Tobacco, <i>Nicotiana suaveolens</i>
5.109 species habitat units for Velvet Daisy-bush, <i>Olearia pannosa</i> subsp. <i>cardiophylla</i>
11.944 species habitat units for Forked Rice-flower, <i>Pimelea hewardiana</i>
12.109 species habitat units for Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
11.797 species habitat units for Brittle Greenhood, <i>Pterostylis truncata</i>
12.109 species habitat units for Fragrant Saltbush, <i>Rhagodia parabolica</i>
4.444 species habitat units for Button Wrinklewort, <i>Rutidosia leptorhynchoides</i>
5.109 species habitat units for Large-headed Fireweed, <i>Senecio macrocarpus</i>
12.109 species habitat units for Cane Spear-grass, <i>Austrostipa breviglumis</i>
12.109 species habitat units for Rye Beetle-grass, <i>Tripogon loliiformis</i>
5.109 species habitat units for Plump Swamp Wallaby-grass, <i>Amphibromus pithogastrus</i>
12.109 species habitat units for Heath Spear-grass, <i>Austrostipa exilis</i>
5.109 species habitat units for Brackish Plains Buttercup, <i>Ranunculus diminutus</i>
12.109 species habitat units for Melbourne Yellow-gum, <i>Eucalyptus leucoxyton</i> subsp. <i>connata</i>
12.109 species habitat units for Matted Flax-lily, <i>Dianella amoena</i>
5.109 species habitat units for Large-flower Crane's-bill, <i>Geranium</i> sp. 1
12.049 species habitat units for Shiny Leionema, <i>Leionema lamprophyllum</i> subsp. <i>obovatum</i>
12.109 species habitat units for Arching Flax-lily, <i>Dianella</i> sp. aff. <i>longifolia</i> (<i>Benambra</i>)

Native vegetation offset report

Habitat units of gain per zone of the proposed offset site

This table provides the habitat units of gain per zone of the offset site. Trading and allocation of units within the **Native Vegetation Credit Register** takes place at the zone.

The species-general offset test is done to determine which species the proposed offset site provides habitat for. The threshold is set at 0.0025 per cent of the mapped habitat value for a species. When the threshold is met or exceeded, species habitat units are generated. If required species habitat units can be generated for all other species mapped at the site. Multiple species units will be generated if the **threshold** is exceeded for multiple species.

The species habitat units for each species in a zone is calculated by the following equation in accordance with the Guidelines:

$$\text{Species habitat units} = \text{extent} \times \text{gain score} \times \text{species landscape factor, where the species landscape factor} = 0.5 + (\text{habitat importance score}/2)$$

The general habitat units in a zone is calculated by the following equation in accordance with the Guidelines:

$$\text{General habitat units} = \text{extent} \times \text{gain score} \times \text{general landscape factor, where the general landscape factor} = 0.5 + (\text{strategic biodiversity value score}/2)$$

Species and general habitat units are alternates and the use or sale of one type of unit will affect the number of other types of units remaining.

Information provided by or on behalf of the applicant				Information calculated by EnSym					
Zone	Type	Gain score	Large tree	Polygon extent	Extent without overlap	SBV	HIS	Habitat units	Attributes
1-S	Patch	0.295	3	2.733	2.733	0.916	0.771	0.771 general habitat units	Port Phillip And Westernport ; Moorabool Shire
							0.800	0.724 species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
							0.480	0.596 species habitat units	10309 Swift Parrot, <i>Lathamus discolor</i>
							0.532	0.617 species habitat units	10504 Speckled Warbler, <i>Chthonicola sagittatus</i>
							0.899	0.764 species habitat units	12922 Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i>
							0.736	0.699 species habitat units	13117 Brown Toadlet, <i>Pseudophryne bibronii</i>

Native vegetation offset report

0.908	0.768 species habitat units	15021 Golden Sun Moth, <i>Synemona plana</i>
0.800	0.724 species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
0.800	0.724 species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
0.736	0.699 species habitat units	501473 Small Golden Moths, <i>Diuris basaltica</i>
0.800	0.724 species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
0.800	0.724 species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
0.800	0.724 species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
0.799	0.724 species habitat units	502317 Velvet Daisy-bush, <i>Olearia pannosa</i> subsp. <i>cardiophylla</i>
0.800	0.724 species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
0.800	0.724 species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
0.796	0.723 species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
0.800	0.724 species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
0.736	0.699 species habitat units	502982 Button Wrinklewort, <i>Rutidosis leptorhynchoides</i>
0.799	0.724 species habitat units	503116 Large-headed Fireweed, <i>Senecio macrocarpus</i>
0.800	0.724 species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
0.800	0.724 species habitat units	503455 Rye Beetle-grass, <i>Tripogon liliiformis</i>

Native vegetation offset report

		0.799	0.724 species habitat units	503624 Plump Swamp Wallaby-grass, <i>Amphibromus pithogastrus</i>		
		0.800	0.724 species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>		
		0.799	0.724 species habitat units	504314 Brackish Plains Buttercup, <i>Ranunculus diminutus</i>		
		0.800	0.724 species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxydon subsp. connata</i>		
		0.800	0.724 species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>		
		0.799	0.724 species habitat units	505342 Large-flower Crane's-bill, <i>Geranium sp. 1</i>		
		0.796	0.723 species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum subsp. obovatum</i>		
		0.800	0.724 species habitat units	505560 Arching Flax-lily, <i>Dianella sp. aff. longifolia (Benambra)</i>		
2-F	Patch	0.639	0.639	0.796	0	0.367
		0.815	0.213 species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>		
		0.897	0.223 species habitat units	12922 Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i>		
		0.915	0.225 species habitat units	15021 Golden Sun Moth, <i>Synemon plana</i>		
		0.815	0.213 species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>		
		0.815	0.213 species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>		
		0.815	0.213 species habitat units	502317 Velvet Daisy-bush, <i>Olearia pannosa subsp. cardiophylla</i>		
		0.815	0.213 species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea var. nivea</i>		

Native vegetation offset report

		0.815	0.213	species habitat units	502929	Fragrant Saltbush, <i>Rhagodia parabolica</i>
		0.815	0.213	species habitat units	503116	Large-headed Fireweed, <i>Senecio macrocarpus</i>
		0.815	0.213	species habitat units	503268	Cane Spear-grass, <i>Austrostipa breviglumis</i>
		0.815	0.213	species habitat units	503455	Rye Beetle-grass, <i>Tripogon loliiformis</i>
		0.815	0.213	species habitat units	503624	Plump Swamp Wallaby-grass, <i>Amphibromus pithogastrus</i>
		0.815	0.213	species habitat units	503984	Heath Spear-grass, <i>Austrostipa exilis</i>
		0.815	0.213	species habitat units	504314	Brackish Plains Buttercup, <i>Ranunculus dimidatus</i>
		0.815	0.213	species habitat units	504484	Melbourne Yellow-gum, <i>Eucalyptus leucoxyloides subsp. connata</i>
		0.815	0.213	species habitat units	505084	Matted Flax-lily, <i>Dianella amoena</i>
		0.815	0.213	species habitat units	505342	Large-flower Crane's-bill, <i>Geranium sp. 1</i>
		0.810	0.212	species habitat units	505478	Shiny Leionema, <i>Leionema lamprophyllum subsp. obovatum</i>
		0.815	0.213	species habitat units	505560	Arching Flax-lily, <i>Dianella sp. aff. longifolia (Benambra)</i>
3-N	Patch	0.262	0.262	0.083	0.730	Port Phillip And Westernport ; Moorabool Shire
		0.808	0.087	species habitat units	505136	Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
		0.891	0.091	species habitat units	12922	Grassland Earless Dragon, <i>Tympanocryptis pinguicollis</i>
		0.911	0.092	species habitat units	15021	Golden Sun Moth, <i>Synemon plana</i>

Native vegetation offset report

0.808	0.087	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
0.790	0.086	species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
0.790	0.086	species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
0.790	0.086	species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
0.808	0.087	species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
0.811	0.087	species habitat units	502317 Velvet Daisy-bush, <i>Olearia pannosa</i> subsp. <i>cardiophylla</i>
0.790	0.086	species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
0.808	0.087	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
0.808	0.087	species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
0.811	0.087	species habitat units	503116 Large-headed Fireweed, <i>Senecio macrocarpus</i>
0.808	0.087	species habitat units	503268 Cane Spear-grass, <i>Auistrostipa breviglumis</i>
0.808	0.087	species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>
0.811	0.087	species habitat units	503624 Plump Swamp Wallaby-grass, <i>Amphibromus pithogastrus</i>
0.808	0.087	species habitat units	503984 Heath Spear-grass, <i>Auistrostipa exilis</i>
0.811	0.087	species habitat units	504314 Brackish Plains Buttercup, <i>Ranunculus diminitus</i>
0.808	0.087	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxyloides</i> subsp. <i>connata</i>

Native vegetation offset report

		0.808	0.087	species habitat units	505084	Matted Flax-lily, <i>Dianella amoena</i>
		0.811	0.087	species habitat units	505342	Large-flower Crane's-bill, <i>Geranium sp. 1</i>
		0.807	0.087	species habitat units	505478	Shiny Leionema, <i>Leionema lamprophyllum subsp. obovatum</i>
		0.808	0.087	species habitat units	505560	Arching Flax-lily, <i>Dianella sp. aff. longifolia (Benambra)</i>
4-T	Patch	1.147	1.147	0.837	0.310	general habitat units ; Moorabool Shire
		0.784	0.301	species habitat units	505136	Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
		0.480	0.250	species habitat units	10309	Swift Parrot, <i>Lathamus discolor</i>
		0.532	0.259	species habitat units	10504	Speckled Warbler, <i>Chthonicola sagittatus</i>
		0.888	0.319	species habitat units	12922	Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i>
		0.745	0.295	species habitat units	13117	Brown Toadlet, <i>Pseudophryne bibronii</i>
		0.901	0.321	species habitat units	15021	Golden Sun Moth, <i>Syneonema plana</i>
		0.784	0.301	species habitat units	500598	Yellow Burr-daisy, <i>Calotis lappulacea</i>
		0.784	0.301	species habitat units	501090	Trailing Hop-bush, <i>Dodonaea procumbens</i>
		0.784	0.301	species habitat units	501518	Western Golden-tip, <i>Goodia medicaginea</i>
		0.784	0.301	species habitat units	501988	Hairy Beard-heath, <i>Leucopogon microphyllus var. pilibundus</i>
		0.784	0.301	species habitat units	502275	Austral Tobacco, <i>Nicotiana suaveolens</i>

Native vegetation offset report

		0.784	0.301	species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
		0.784	0.301	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
		0.784	0.301	species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
		0.784	0.301	species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
		0.784	0.301	species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
		0.784	0.301	species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>
		0.784	0.301	species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
		0.784	0.301	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon</i> subsp. <i>connata</i>
		0.784	0.301	species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
		0.776	0.300	species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum</i> subsp. <i>obovatum</i>
		0.784	0.301	species habitat units	505560 Arching Flax-lily, <i>Dianella</i> sp. aff. <i>longifolia</i> (<i>Benambra</i>)
5-N	Patch	0.367	0	0.960 0.960 0.730	Port Phillip And Westernport ; Moorabool Shire
		0.798	0.317	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
		0.890	0.333	species habitat units	12922 Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i>
		0.910	0.337	species habitat units	15021 Golden Sun Moth, <i>Synemon plana</i>
		0.798	0.317	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>

Native vegetation offset report

0.793	0.316 species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
0.810	0.319 species habitat units	501473 Small Golden Moths, <i>Diuris basaltica</i>
0.793	0.316 species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
0.810	0.319 species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
0.798	0.317 species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
0.803	0.318 species habitat units	502317 Velvet Daisy-bush, <i>Olearia pannosa</i> subsp. <i>cardiophylla</i>
0.793	0.316 species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
0.798	0.317 species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
0.810	0.319 species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
0.798	0.317 species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
0.803	0.318 species habitat units	503116 Large-headed Fireweed, <i>Senecio macrocarpus</i>
0.798	0.317 species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
0.798	0.317 species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>
0.803	0.318 species habitat units	503624 Plump Swamp Wallaby-grass, <i>Amphibromus pithogastrus</i>
0.798	0.317 species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
0.803	0.318 species habitat units	504314 Brackish Plains Buttercup, <i>Ranunculus dimidatus</i>

Native vegetation offset report

				0.798	0.317	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon subsp. connata</i>
				0.798	0.317	species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
				0.803	0.318	species habitat units	505342 Large-flower Crane's-bill, <i>Geranium sp. 1</i>
				0.793	0.316	species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum subsp. obovatum</i>
				0.798	0.317	species habitat units	505560 Arching Flax-lily, <i>Dianella sp. aff. longifolia (Benambra)</i>
6-S	Patch	0.295	1	0.337	0.337	0.856	Port Phillip And Westernport ; Moorabool Shire
				0.787	0.089	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
				0.480	0.073	species habitat units	10309 Swift Parrot, <i>Lathamus discolor</i>
				0.535	0.076	species habitat units	10504 Speckled Warbler, <i>Chthonicola sagittatus</i>
				0.745	0.086	species habitat units	13117 Brown Toadlet, <i>Pseudophryne bibronii</i>
				0.906	0.094	species habitat units	15021 Golden Sun Moth, <i>Synemon plana</i>
				0.787	0.089	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
				0.787	0.089	species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
				0.787	0.089	species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
				0.787	0.089	species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus var. pilibundus</i>
				0.787	0.089	species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>

Native vegetation offset report

		0.787	0.089	species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
		0.787	0.089	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
		0.787	0.089	species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
		0.787	0.089	species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
		0.787	0.089	species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
		0.787	0.089	species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliformis</i>
		0.787	0.089	species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
		0.787	0.089	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon</i> subsp. <i>connata</i>
		0.787	0.089	species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
		0.791	0.089	species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum</i> subsp. <i>obovatum</i>
		0.787	0.089	species habitat units	505560 Arching Flax-lily, <i>Dianella</i> sp. aff. <i>longifolia</i> (<i>Benambra</i>)
7-S	Patch	1.628	0.439	general habitat units	Port Phillip And Westernport ; Moorabool Shire
		0.779	0.426	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
		0.480	0.355	species habitat units	10309 Swift Parrot, <i>Lathamus discolor</i>
		0.534	0.368	species habitat units	10504 Speckled Warbler, <i>Chthonicola sagittatus</i>
		0.884	0.452	species habitat units	12922 Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i>
		0.777	0.426	species habitat units	13117 Brown Toadlet, <i>Pseudophryne bibronii</i>

Native vegetation offset report

0.901	0.456 species habitat units	15021 Golden Sun Moth, <i>Synemon plana</i>
0.779	0.426 species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
0.779	0.426 species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
0.779	0.426 species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
0.779	0.426 species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
0.779	0.426 species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
0.779	0.426 species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
0.779	0.426 species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
0.778	0.426 species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
0.779	0.426 species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
0.779	0.426 species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
0.779	0.426 species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>
0.779	0.426 species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
0.779	0.426 species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon</i> subsp. <i>connata</i>
0.779	0.426 species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
0.780	0.427 species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum</i> subsp. <i>obovatum</i>

Native vegetation offset report

8-M	Patch	0.367	0	0.482	0.482	0.866	0.779	0.426	species habitat units	505560 Arching Flax-lily, <i>Dianella</i> sp. aff. <i>longifolia</i> (Benambra)
										Port Phillip And Westernport ; Moorabool Shire
							0.778	0.157	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
							0.510	0.134	species habitat units	10504 Speckled Warbler, <i>Chthonicola sagittatus</i>
							0.906	0.169	species habitat units	15021 Golden Sun Moth, <i>Synemon plana</i>
							0.778	0.157	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
							0.778	0.157	species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
							0.778	0.157	species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
							0.777	0.157	species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
							0.778	0.157	species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
							0.778	0.157	species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
							0.778	0.157	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
							0.777	0.157	species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
							0.778	0.157	species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
						0.778	0.157	species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>	
						0.778	0.157	species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>	

Native vegetation offset report

		0.778	0.157	species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
		0.778	0.157	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon subsp. connata</i>
		0.778	0.157	species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
		0.804	0.160	species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum subsp. obovatum</i>
		0.778	0.157	species habitat units	505560 Arching Flax-lily, <i>Dianella sp. aff. longifolia (Benambra)</i>
9-M	Patch	0.964	0.367	0	Port Phillip And Westernport ; Moorabool Shire
		0.773	0.314	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
		0.480	0.262	species habitat units	10309 Swift Parrot, <i>Lathamus discolor</i>
		0.506	0.267	species habitat units	10504 Speckled Warbler, <i>Chthonicola sagittatus</i>
		0.750	0.310	species habitat units	13117 Brown Toadlet, <i>Pseudophryne bibronii</i>
		0.902	0.337	species habitat units	15021 Golden Sun Moth, <i>Synemona plana</i>
		0.773	0.314	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
		0.773	0.314	species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
		0.773	0.314	species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
		0.773	0.314	species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus var. pilibundus</i>
		0.773	0.314	species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>

Native vegetation offset report

		0.773	0.314	species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
		0.773	0.314	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
		0.772	0.314	species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
		0.773	0.314	species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
		0.773	0.314	species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
		0.773	0.314	species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>
		0.773	0.314	species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
		0.773	0.314	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon</i> subsp. <i>connata</i>
		0.773	0.314	species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
		0.778	0.315	species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum</i> subsp. <i>obovatum</i>
		0.773	0.314	species habitat units	505560 Arching Flax-lily, <i>Dianella</i> sp. aff. <i>longifolia</i> (<i>Benambra</i>)
10-N	Patch	0.182	0.182	0.059	Port Phillip And Westernport ; Moorabool Shire
		0.788	0.060	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
		0.788	0.060	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
		0.788	0.060	species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
		0.788	0.060	species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>

Native vegetation offset report

		0.788	0.060	species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
		0.788	0.060	species habitat units	502275 Austral Tobacco, <i>Nicotiana</i> <i>suaveolens</i>
		0.788	0.060	species habitat units	502522 Forked Rice-flower, <i>Pimelea</i> <i>hewardiana</i>
		0.788	0.060	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
		0.780	0.059	species habitat units	502821 Brittle Greenhood, <i>Pterostylis</i> <i>truncata</i>
		0.788	0.060	species habitat units	502929 Fragrant Saltbush, <i>Rhagodia</i> <i>parabolica</i>
		0.788	0.060	species habitat units	503268 Cane Spear-grass, <i>Austrostipa</i> <i>breviglumis</i>
		0.788	0.060	species habitat units	503455 Rye Beetle-grass, <i>Tripogon</i> <i>loliiformis</i>
		0.788	0.060	species habitat units	503984 Heath Spear-grass, <i>Austrostipa</i> <i>exilis</i>
		0.788	0.060	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon</i> subsp. <i>connata</i>
		0.788	0.060	species habitat units	505084 Matted Flax-lily, <i>Dianella</i> <i>amoena</i>
		0.788	0.060	species habitat units	505478 Shiny Leionema, <i>Leionema</i> <i>lamprophyllum</i> subsp. <i>obovatum</i>
		0.788	0.060	species habitat units	505560 Arching Flax-lily, <i>Dianella</i> sp. <i>aff. longifolia</i> (Benambra)
12-N	Patch	0.041	0.041	0.828	0.014 general habitat units Port Phillip And Westernport ; Moorabool Shire
		0.794	0.014	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia</i> <i>rostriformis</i>
		0.510	0.011	species habitat units	10504 Speckled Warbler, <i>Chthonicola</i> <i>sagittatus</i>

Native vegetation offset report

0.910	0.015	species habitat units	15021 Golden Sun Moth, <i>Synemona plana</i>
0.794	0.014	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
0.794	0.014	species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
0.794	0.014	species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
0.800	0.014	species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
0.794	0.014	species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
0.794	0.014	species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
0.794	0.014	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
0.800	0.014	species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
0.794	0.014	species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
0.794	0.014	species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
0.794	0.014	species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>
0.794	0.014	species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
0.794	0.014	species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon</i> subsp. <i>connata</i>
0.794	0.014	species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
0.794	0.014	species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum</i> subsp. <i>obovatum</i>

Native vegetation offset report

13-R	Patch	0.295	1	0.794	0.014	species habitat units	505560 Arching Flax-lily, <i>Dianella</i> sp. aff. <i>longifolia</i> (Benambra)
				21.938	21.938	0.936	Port Phillip And Westernport ; Moorabool Shire
				0.745	5.638	species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
				0.721	5.558	species habitat units	507580 Werribee Blue-box, <i>Eucalyptus baueriana</i> subsp. <i>thlassina</i>
				0.477	4.773	species habitat units	10309 Swift Parrot, <i>Lathamus discolor</i>
				0.519	4.908	species habitat units	10504 Speckled Warbler, <i>Chthonicola sagittatus</i>
				0.873	6.050	species habitat units	12922 Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i>
				0.799	5.811	species habitat units	13117 Brown Toadlet, <i>Pseudophryne bibronii</i>
				0.893	6.116	species habitat units	15021 Golden Sun Moth, <i>Synemon plana</i>
				0.746	5.641	species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
				0.755	5.669	species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>
				0.745	5.638	species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
				0.745	5.638	species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
				0.745	5.638	species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
0.747	5.643	species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>				
0.745	5.638	species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>				

Native vegetation offset report

		0.745	5.638 species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
		0.745	5.638 species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
		0.745	5.638 species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
		0.745	5.638 species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliiformis</i>
		0.745	5.638 species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
		0.745	5.638 species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxydon subsp. connata</i>
		0.745	5.638 species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
		0.730	5.590 species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum subsp. obovatum</i>
		0.745	5.638 species habitat units	505560 Arching Flax-lily, <i>Dianella sp. aff. longifolia (Benambra)</i>
14-M	Patch	11.555	3.923 general habitat units	Port Phillip And Westernport ; Moorabool Shire
		0.779	3.774 species habitat units	505136 Bacchus Marsh Wattle, <i>Acacia rostriformis</i>
		0.522	3.228 species habitat units	10504 Speckled Warbler, <i>Chthonicola sagittatus</i>
		0.877	3.981 species habitat units	12922 Grassland Earless Dragon, <i>Tympanocryptis pinguicolla</i>
		0.899	4.028 species habitat units	15021 Golden Sun Moth, <i>Synemon plana</i>
		0.780	3.776 species habitat units	500598 Yellow Burr-daisy, <i>Calotis lappulacea</i>
		0.797	3.813 species habitat units	501090 Trailing Hop-bush, <i>Dodonaea procumbens</i>

Native vegetation offset report

0.763	3.740 species habitat units	501473 Small Golden Moths, <i>Diuris basaltica</i>
0.797	3.813 species habitat units	501518 Western Golden-tip, <i>Goodia medicaginea</i>
0.798	3.815 species habitat units	501988 Hairy Beard-heath, <i>Leucopogon microphyllus</i> var. <i>pilibundus</i>
0.776	3.768 species habitat units	502275 Austral Tobacco, <i>Nicotiana suaveolens</i>
0.775	3.767 species habitat units	502317 Velvet Daisy-bush, <i>Olearia pannosa</i> subsp. <i>cardiophylla</i>
0.797	3.813 species habitat units	502522 Forked Rice-flower, <i>Pimelea hewardiana</i>
0.776	3.768 species habitat units	502746 Snowy Mint-bush, <i>Prostanthera nivea</i> var. <i>nivea</i>
0.771	3.757 species habitat units	502821 Brittle Greenhood, <i>Pterostylis truncata</i>
0.776	3.768 species habitat units	502929 Fragrant Saltbush, <i>Rhagodia parabolica</i>
0.765	3.745 species habitat units	502982 Button Wrinklewort, <i>Rutidosis leptorhynchoides</i>
0.775	3.767 species habitat units	503116 Large-headed Fireweed, <i>Senecio macrocarpus</i>
0.776	3.768 species habitat units	503268 Cane Spear-grass, <i>Austrostipa breviglumis</i>
0.776	3.768 species habitat units	503455 Rye Beetle-grass, <i>Tripogon loliformis</i>
0.775	3.767 species habitat units	503624 Plump Swamp Wallaby-grass, <i>Amphibromus pithogastrus</i>
0.776	3.768 species habitat units	503984 Heath Spear-grass, <i>Austrostipa exilis</i>
0.775	3.767 species habitat units	504314 Brackish Plains Buttercup, <i>Ranunculus dimidatus</i>

Native vegetation offset report

0.776	3.768 species habitat units	504484 Melbourne Yellow-gum, <i>Eucalyptus leucoxylon subsp. connata</i>
0.776	3.768 species habitat units	505084 Matted Flax-lily, <i>Dianella amoena</i>
0.775	3.767 species habitat units	505342 Large-flower Crane's-bill, <i>Geranium sp. 1</i>
0.771	3.758 species habitat units	505478 Shiny Leionema, <i>Leionema lamprophyllum subsp. obovatum</i>
0.776	3.768 species habitat units	505560 Arching Flax-lily, <i>Dianella sp. aff. longifolia (Benambra)</i>

Next steps

Offset sites must meet eligibility criteria as outlined in the *Guidelines for the removal, destruction or lopping of native vegetation* and the *Native vegetation gain scoring manual, version 2* available on the DELWP website, and any other relevant requirements. Eligible offset sites that are intended to be banked or sold as credits must be registered on the Native Vegetation Credit Register (NVCR). A gain scoring assessment must be done before any offset can be registered on the NVCR. All proposed offset sites must be secured by a relevant security agreement that includes an offset management plan.

© The State of Victoria Department of Environment, Land, Water and Planning Melbourne 2018

This work is licensed under a Creative Commons Attribution 4.0 International licence. You are free to re-use the work under that licence, on the condition that you credit the State of Victoria as author. The licence does not apply to any images, photographs or branding, including the Victorian Coat of Arms, the Victorian Government logo and the Department of Environment, Land, Water and Planning logo. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/34.0/au/deed.en>

Authorised by the Victorian Government, 8 Nicholson Street, East Melbourne.

For more information contact the DELWP Customer Service Centre 136 186

Disclaimer

This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.


Appendix 1 – Images of marked native vegetation

1. Aerial photograph showing marked native vegetation




Native vegetation offset report

2. Strategic biodiversity value map




3. Habitat importance maps

Bacchus Marsh Wattle
Acacia rostriformis
505136




Werribee Blue-box
Eucalyptus baueriana subsp. thalassina
507580




Native vegetation offset report


Swift Parrot
Lathamus discolor
10309




Speckled Warbler
Chthonicola sagittatus
10504



Grassland Earless Dragon
Tympanocryptis pinguicolla
12922





Brown Toadlet
Pseudophryne bibronii
13117





Native vegetation offset report



Golden Sun Moth
Synemon plana
15021





Yellow Burr-daisy
Calotis lappulacea
500598



Trailing Hop-bush
Dodonaea procumbens
501090




Small Golden Moths
Diuris basaltica
501473




Native vegetation offset report


Western Golden-tip
Goodia medicaginea
501518




Hairy Beard-heath
Leucopogon microphyllus var. *pilibundus*
501988



Austral Tobacco
Nicotiana suaveolens
502275




Velvet Daisy-bush
Olearia pannosa subsp. *cardiophylla*
502317




Native vegetation offset report


Forked Rice-flower
Pimelea hewardiana
502522




Snowy Mint-bush
Prostanthera nivea var. *nivea*
502746



Brittle Greenhood
Pterostylis truncata
502821




Fragrant Saltbush
Rhagodia parabolica
502929




Native vegetation offset report


Button Wrinklewort
Rutidosia leptorhynchoides
502982




Large-headed Fireweed
Senecio macrocarpus
503116



Cane Spear-grass
Austrostipa breviglumis
503268




Rye Beetle-grass
Tripogon loliiformis
503455




Native vegetation offset report


Plump Swamp Wallaby-grass
Amphibromus pithogastrus
503624




Heath Spear-grass
Austrostipa exilis
503984



Brackish Plains Buttercup
Ranunculus diminutus
504314




Melbourne Yellow-gum
Eucalyptus leucoxylon subsp. connata
504484




Native vegetation offset report


Matted Flax-lily
Dianella amoena
505084




Large-flower Crane's-bill
Geranium sp. 1
505342



Shiny Leionema
Leionema lamprophyllum subsp. obovatum
505478



Arching Flax-lily
Dianella sp. aff. longifolia (Benambra)
505560



GLOSSARY

Alternate offset types	Offset types within a zone are alternates. The use of one offset type will result in the proportional reduction of all other offset types within the zone. Refer to <i>Native vegetation offset sites</i> fact sheet available on the DELWP website for more information.
Gain score	This is the site-assessed gain score for the native vegetation based on the agreed management and security commitments. Each zone in the proposed offset site is assigned a gain score according to the gain scoring assessment. The score is divided by 100 to give a number between 0 and 1.
General habitat units of gain	The general habitat units quantify the overall contribution that the protection and management of native vegetation at the offset site makes to Victoria's biodiversity. The general habitat units are calculated as follows: $\text{General habitat units} = \text{extent} \times \text{gain score} \times \text{general landscape factor}$
General landscape factor	The general landscape factor is the adjusted strategic biodiversity value (SBV) score. The SBV score is adjusted so that site-based biodiversity information has more influence on the number of units.
General offset attributes	The attributes of a general offset includes the location (Catchment Management Authority and Municipal District), strategic biodiversity value score and the number of large trees protected.
Offset type	There are two types of offsets, general offsets and species offsets. All offset sites include general offsets. Sites that are mapped as habitat for rare or threatened species can also include species offsets for the mapped species.
Species offset attributes	The attributes of a species offset is the mapped habitat for the species and the number of large trees protected.
Species habitat units of gain	The species habitat units quantify the overall contribution that the protection and management of native vegetation at an offset site makes to the habitat of the relevant rare or threatened species. Species habitat units are calculated for each species in the zone where the result of the threshold test is greater than 0.0025 per cent. Species units are calculated as follows: $\begin{aligned} \text{Species habitat units}_{\text{species } x} \\ = \text{extent} \times \text{gain score} \times \text{species landscape factor}_{\text{species } x} \end{aligned}$